arXiv:q-alg/9712012v1 3 Dec 1997

PERFECT CRYSTALS OF $U_q(G_2^{(1)})$

Shigenori Yamane

Department of Mathematical Science,Faculty of Engineering Science,
Osaka University, Toyonaka, Osaka 560, Japan

yamane@sigmath.es.osaka-u.ac.jp

1 Introduction

The theory of *crystal base* was introduced by Kashiwara in [3]. He proved the existence and uniqueness of this base for a quantized universal enveloping algebras $U_q(\mathfrak{g})$, where $\mathfrak{g}$ is an arbitrary Kac-Moody Lie algebra with symmetrizable Cartan matrix. The theory of crystal base provides a powerful combinatorial tool for studying the structure of the integrable highest weight modules and the decomposition of tensor products of them.

Let $\mathfrak{g}$ be an arbitrary Kac-Moody Lie algebra with a symmetrizable Cartan matrix. Let I be its index set. Denote the set of simple roots of $\mathfrak{g}$ by $\{\alpha_i \mid i \in I\}$. Let P be the weight lattice of $\mathfrak{g}$. Take an integrable $U_q(\mathfrak{g})$-module M and let $\tilde{f}_i$, $\tilde{e}_i$ be the Kashiwara operators (cf. §3.1.2) on M. Let $\mathbf{A}$ be the subring of $\mathbf{Q}(q)$ consisting of $f \in \mathbf{Q}(q)$ that is regular at $q = 0$. A crystal lattice L of an integrable $U_q(\mathfrak{g})$-module M is free $\mathbf{A}$-submodule of M such that $M \cong \mathbf{Q}(q) \otimes_{\mathbf{A}} L$, $L = \bigoplus_{\lambda \in P} L_\lambda$, where $L_\lambda = L \cap M_\lambda$ and $\tilde{e}_i L \subset L$, $\tilde{f}_i L \subset L$. A crystal base of the integrable $U_q(\mathfrak{g})$-module M is a pair (L, B) such that (i) L is a cyrstal lattice of M, (ii) B is a $\mathbf{Q}$-base of L/qL, (iii) $B = \sqcup_{\lambda \in P} B_\lambda$ where $B_\lambda = B \cap (L_\lambda / qL_\lambda)$, (iv)$\tilde{e}_i B \subset B \sqcup \{0\}$, and (v) for $b, b' \in B$, $b' = \tilde{f}_i b$ if and only if $b = \tilde{e}_i b'$ for $i \in I$. We sometimes replace condition (ii) by : $B_{ps} = B' \sqcup (-B')$ where B' is a $\mathbf{Q}$-base of L/qL. In this case, w e call (L, B_{ps}) a *crystal pseudo base.* The quotient $B_{ps}/\{\pm 1\}$ is called an *associated crystal* of (L, B_{ps}).

In [6], Kashiwara and Nakashima gave an explicit construction of crystal bases for all finite demensional irreducible modules over $U_q(\mathfrak{g})$, where $\mathfrak{g}$ is of type A_n, B_n, C_n, D_n, in terms of generalized Young tableaux. Kang and Misra gave a construction of crystal bases of $U_q(G_2)$-module by a method similar to Kashiwara and Nakashima's in [7].

The notion of *perfect crystals* was introduced in [8] and [9] in order to compute the one-point functions of vertex model in 2 dimensional lattice statistical models. If a $U_q(\mathfrak{g})$-module M has perfect crystal base, we have the notion of paths. Using paths, we can compute one-point functions of vertex models. We define classical part of the weight lattice by $P_{cl} = \sum \mathbf{Z}\Lambda_i$. A P_{cl}-weighted crystal is called a *classical crystal* and P-weighted crystal is called an *affine crystal.* Let c be the central element. Further we define $\varepsilon_i(b)$, $\varphi_i(b)$ by $\max\{k \geq 0 \mid \tilde{e}_i^k b \neq 0\}$, $\max\{k \geq 0 \mid \tilde{f}_i^k b \neq 0\}$ respectively. We define $\varepsilon(b)$, $\varphi(b)$ by $\sum_i \varepsilon_i(b)\Lambda_i$, $\sum_i \varphi_i(b)\Lambda_i$ respectively. A classical crystal B is called a *perfect crystal of level l* if B satisfies: (i) $B \otimes B$ is connected, (ii) there exists $\lambda_0 \in P_{cl}$ such that $\mathrm{wt}(B) \in \lambda_0 + \sum_{i \neq 0} \mathbf{Z}_{\leq 0} cl(\alpha_i)$ and that $\sharp(B_{\lambda_0}) = 1$, (iii) there is a finite dimensional $U'_q(\mathfrak{g})$-module with a crystal pseudo base (L, B_{ps}) such that $B \cong B_{ps}/\pm 1$, (iv) for any $b \in B$, we have $\langle c, \varepsilon(b) \rangle \geq l$, and (v) the maps ε, φ : $B_l = \{b \in B \,|\, \langle c, \varepsilon(b) \rangle = l\} \to (P_{cl}^+)_l = \{\lambda \in \sum \mathbf{Z}_{\geq 0}\Lambda_i \,|\, \langle c, \lambda \rangle = l\}$ are bijective.

In [8], they give examples of perfect crystals of arbitrary levels for algebras of the following types: $A_n^{(1)}$, $B_n^{(1)}$, $C_n^{(1)}$, $D_n^{(1)}$, $A_{2n}^{(2)}$, $A_{2n-1}^{(2)}$, $D_{n+1}^{(2)}$. Up to now we do not know simple criterion for perfectness of crystals. Further, perfect crystals with arbirary level for $E_n^{(1)}$, $F_4^{(1)}$, $G_2^{(1)}$, $E_6^{(2)}$ are not known. In this paper, we give a series of perfect crystals of $U_q(G_2^{(1)})$, following method in [8].

Let $U'_q(G_2^{(1)})$ be the subalgebra of $U_q(G_2^{(1)})$, generated by e_i, f_i, q^h ($h \in P_{cl}^* = \sum_{i \in \{0,1,2\}} \mathbf{Z}h_i$). We proceed in the following way.

1. As in [8], for a given level l, our strategy is to emply the fusion procedure in order to construct finite dimensional irreducible module V_l of $U'_q(G_2^{(1)})$ with a crystal pseudo base such that its associated crystal is perfect of level l. As a starting point, we take the following data. Let V be the direct sum of the 14-dimensional module with highest weight Λ_1 and the trivial module over $U_q(G_2)$. The module V has a crystal base which is characterized by a polarization on V. A polarization on $U_q(\mathfrak{g})$-module M

is the symmetric bilinear such that for $u, v \in M$ (i)$(q^h u, v) = (u, q^h v)$, (ii) $(e_i u, v) = (u, q_i^{-1} t_i^{-1} f_i v)$, (iii) $(f_i u, v) = (u, q_i^{-1} t_i e_i v)$ and (iv) positive definite. Let B^1 be the associated crystal of $U_q(G_2)$. We define the actions of e_0, f_0, and q^{h_0} on V to make it an irreducible module for $U_q'(G_2^{(1)})$. Then we show that there exists a polarization for $U_q'(G_2^{(1)})$. We see that B^1 is a perfect crystal of level 1 by direct calculation. Using the global base [5], we compute the R-matrix for V explicitly. By the fusion construction, we obtain a certain finite-dimensional submodule V_l of $V^{\otimes l}$. The existence of polarization on V_l ensures that the submodule V_l has a crystal pseudo base. The associated crystal of V_l is isomorphic to a crystal of $U_q(G_2)$ and a crystal of $U_q(A_2)$ as crystals of $U_q(G_2)$ and $U_q(A_2)$ respectively. These isomorphisms show that the associated crystal is a crystal of $U_q'(G_2^{(1)})$. It is a perfect crystal of level l. This is shown in the section 4.

2. In section 5, we construct a perfect crystal B^l of $U_q(G_2^{(1)})$ in a combinatorial way.

   The Dynkin diagram of $G_2^{(1)}$ is . Here, let $J_0 = \{1, 2\}$ and $J_2 = \{1, 0\}$ be the index set of G_2 and A_2, respectively. We define $\imath_i : J_i \longrightarrow I$ by $\imath_i(j) = j$ $(i = 0, 2)$. We define crystal $\mathcal{G}^l$ to be

   $$\mathcal{G}^l = \bigoplus_{n=0}^{l} B^{G_2}(n\Lambda_1) \text{ (as crystals for } U_q(G_2) \text{).}$$

   The affine crystal b^l for $U_q(G_2^{(1)})$ is constructed with $\mathcal{G}^l$ and $\tilde{f}_0$. To define $\tilde{f}_0$ on $\mathcal{G}^l$, we use crystals of $U_q(A_2)$. Here, we define $\mathcal{A}$ by

   $$\begin{aligned} \mathcal{A}_i &= \bigoplus_{i \le j_1, j_0 \le l-i} B^{A_2}(j_1\Lambda_1 + j_0\Lambda_0) \text{ (as crystals for } U_q(A_2)), \\ \mathcal{A} &= \bigoplus_{i=0}^{\left[\frac{l}{2}\right]} \mathcal{A}_i. \end{aligned}$$

   We define operators $E_{\mathcal{A}}$, $F_{\mathcal{A}}$ on $\mathcal{A}$ such that for $b, b' \in \mathcal{A}$, (C1) $E_{\mathcal{A}} b = b'$, if and only if $F_{\mathcal{A}} b' = b$, (C2) $E_{\mathcal{A}} \tilde{f}_0^{\mathcal{A}}(b) = \tilde{f}_0^{\mathcal{A}} E_{\mathcal{A}}(b)$, (C3) $\max\{m \mid F_{\mathcal{A}}^m b \neq 0\} - \max\{m' \mid E_{\mathcal{A}}^{m'} b \neq 0\} = -2\mathrm{wt}_1^{\mathcal{A}}(b) - \mathrm{wt}_0^{\mathcal{A}}(b)$. Operators $E_{\mathcal{A}}$ and $F_{\mathcal{A}}$ are counterpart of $\tilde{e}_2^{\mathcal{G}}$, $\tilde{f}_2^{\mathcal{G}}$ on $\mathcal{G}^l$. The crystal base $\mathcal{A}$ has an involution $\mathrm{C}_{\mathcal{A}}$ such that

   $$\mathrm{C}_{\mathcal{A}}\left(\left(\tilde{f}_0^{\mathcal{A}}\right)^r \left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}_{(k,j)}^{l,i}\right) = \left(\tilde{f}_0^{\mathcal{A}}\right)^{j+q-2p-r} \left(\tilde{f}_1^{\mathcal{A}}\right)^{k+j-q} \left(\tilde{f}_0^{\mathcal{A}}\right)^{k-q+p} \overline{b}_{(j,k)}^{l,i},$$

   where $0 \le i \le \left[\frac{l}{2}\right]$, $i \le k, j \le l - i$, $\overline{b}_{(k,j)}^{l,i}$ is the highest weight element on $\mathcal{A}_i$ with weight $j\Lambda_1 + k\Lambda_0$, $0 \le p \le j$, $p \le q \le p + k$, $0 \le r \le j + q - 2p$. By calculation, we see $\sharp\mathcal{A} = \sharp\mathcal{G}^l$. In view of this, we construct one-to-one map $\Phi : \mathcal{A} \rightarrow \mathcal{G}^l$ such that for $b \in \mathcal{A}$ (E1) $\tilde{e}_1^{\mathcal{G}}\Phi(b) = \Phi\left(\tilde{e}_1^{\mathcal{A}} b\right)$, (E2) $\tilde{f}_2^{\mathcal{G}}\Phi(b) = \Phi\left(F_{\mathcal{A}} b\right)$, (E3) $\mathrm{wt}_1^{\mathcal{G}}\left(\Phi(b)\right) = \mathrm{wt}_1^{\mathcal{A}}(b)$, (E4) $\mathrm{wt}_2^{\mathcal{G}}\left(\Phi(b)\right) = -2\mathrm{wt}_1^{\mathcal{A}}(b) - \mathrm{wt}_0^{\mathcal{A}}(b)$, (E5) $\tilde{e}_0^{\mathcal{G}}\Phi(b) = 0$ (resp. $\tilde{f}_0^{\mathcal{G}}\Phi(b) = 0$) if and only if $\tilde{e}_0^{\mathcal{A}}(b) = 0$ (resp. $\tilde{f}_0^{\mathcal{A}}(b) = 0$), for $b \in \mathcal{A}$. Then we define $\tilde{f}_0$ on $\mathcal{G}^l$ by $\Phi\tilde{f}_0^{\mathcal{A}}\Phi^{-1}$. By weight consideration, (C2), involution, etc., we see that the action of $\tilde{f}_0$ is unique. Therefore we have crystal base of $U_q(G_2^{(1)})$.

3. We show that perfectness are satisfied by direct calculation (§5.4,§5.5).

Graphs of the perfect crystal of level 1 and level 2 are given by Figure 1 and 9 respectively. In Figure 9, for simplicity, we omit arrows of $E_{\mathcal{A}}(b)$ such that $\varepsilon_0^{\mathcal{A}}(b) \neq 0$. By definition, we have $B^2 = B^{G_2}(2\Lambda_1) \oplus B^{G_2}(\Lambda_1) \oplus B^{G_2}(\phi)$ and $\mathcal{A}_0 = \oplus_{0 \le j_1, j_0 \le 2} B^{A_2}(j_1\Lambda_1 + j_0\Lambda_0)$, $\mathcal{A}_1 = B^{A_2}(\Lambda_1 + \Lambda_0)$, $\mathcal{A} = \mathcal{A}_0 \oplus \mathcal{A}_1$. In this graph, the element $b \in B^2$ from which we draw arrows of $E_{\mathcal{A}}$ satisfies $\Phi(b) \in B^{G_2}(2\Lambda_1)$, $\Phi\left(\tilde{f}_0^{\mathcal{A}} b\right) \notin B^{G_2}(\Lambda_1)$ and $\tilde{e}_0^{\mathcal{A}} b = 0$. For $b \in \mathcal{A}$ such that $\Phi(b) \in B^{G_2}(\Lambda_1) \oplus B^{G_2}(\phi)$, the action of $E_{\mathcal{A}}$ is same as the case of level 1. Using commutativity (C2), we obtain the completed graph of the perfect crystal of level 2. We obtain the perfect crystal B^1 of level 1 by restricting B^2 to $B^{G_2}(\Lambda_1) \oplus B^{G_2}(\phi)$. From these examples, we expect that we can obtain the perfect crystal B^{l+1} by extending B^l.

To find perfect crystals of level l, we wrote a C program which judges whether a given representation has perfect crystal on a combinatorial level. In this program, we construct crystal bases B using a Lakshmibai-Seshadri path [11]. A Lakshmibai-Seshadri path is a piecewise linear paths $\pi : [0,1] \to (\sum \mathbf{Z}\Lambda_i) \otimes_Z \mathbf{R}$. Operators e_α and f_α on the path are modification by simple reflection σ with respect to the simple root α. When translated into the language of crystal base, a path π is a base and operators e_{α_i} and f_{α_i} are Kashiwara operators $\tilde{e}_i$ and $\tilde{f}_i$. Let $\delta = \sum a_i\alpha_i$ be the null root. We define the operator $\tilde{f}_0 b$ ($b \in B$), by searching bases with weight $\mathrm{wt}(b) + \sum_{i\neq 0} a_i\alpha_i$ and taking into account the conditions for perfectness. The conditions are $\langle c, \varphi(b)\rangle \geq l$, and the map $\varphi : B_l = \{b \in B \mid \langle c, \varphi(b)\rangle = l\} \to \left(P^+_{cl}\right)_l = \{\lambda \in \sum \mathbf{Z}_{\geq 0}\Lambda_i \mid \langle c, \lambda\rangle = l\}$ are bijective (c is the element of the center). If we find suitable action of $\tilde{f}_0$, B is perfect crystal on a combinatorial level.

Using this program we find that $\oplus_{k=0}^{l} B(k\Lambda_1)$ is a candidate of perfect crystals of level l also for $U_q(F_4^{(1)})$, $U_q(E_6^{(1)})$, etc, on a combinatorial level. We hope to report on these in a future publication.

The author would like to acknowledge professor Etsuro Date and T. Nakashima for their helpful advices.



Figure 1: Perfect crystal of level 1

2 Results

Denote the index set of the simple roots of $G_2^{(1)}$ by $I = \{0,1,2\}$. We can take $J_0 = \{1,2\}$ and $J_2 = \{1,0\}$ as the index set of G_2 and A_2, respectively. We define $\imath_i : J_i \to I$ by $\imath_i(j) = j$ ($i = 0,2$).

Proposition 2.1 For any integer $l \geq 1$, there exists the crystal B^l for $G_2^{(1)}$ such that

$$\imath_0^*(B^l) \cong \bigoplus_{k=0}^{l} B^{G_2}(k\Lambda_1) \text{ (as crystals of } U_q(G_2)),$$

and

$$i_2^*(B^l) \cong \bigoplus_{i=0}^{[\frac{l}{2}]} \bigoplus_{i \le j_1, j_0 \le l-i} B^{A_2}(j_1\Lambda_1 + j_0\Lambda_0) \text{ (as crystals of } U_q(A_2)).$$

Theorem 2.2 *The crystal B^l is a perfect crystal of level l for $U_q(G_2^{(1)})$.*

3 Preliminaries

3.1 Crystal base

We recall definitions of quantum enveloping algebras $U_q(A_2)$, $U_q(G_2)$, $U_q(G_2^{(1)})$ and crystal bases of $U_q(A_2)$, $U_q(G_2)$.

3.1.1 Definition of $U_q(\mathfrak{g})$

Let $\mathfrak{g}$ be a semi simple (resp. affine) Lie algebra generated by e_i, f_i $(i \in I = \{1,\ldots,n\}(resp.\{0,1,\ldots,n\}))$ and $\mathfrak{h}$ the Cartan subalgebra over $\mathbf{Q}$. Let $\{\alpha_i \mid i \in I\} \subset \mathfrak{h}^*$ and $\{h_i \mid i \in I\} \subset \mathfrak{h}$ denote the simple roots and simple coroots, respectively. We denote a non-degenerate invariant symmetric bilinear form on $\mathfrak{h}^*$ by $(,)$. This bilinear form $(,)$ satisfies $(\alpha_i,\alpha_i) \in \mathbf{Z}_{>0}$. Let P be the weight lattice and P^* be the dual lattice.

The quantized universal enveloping algebra $U_q(\mathfrak{g})$ is the $\mathbf{Q}(q)$-algebra generated by the symbols e_i, f_i $(i \in I)$ and q^h $(h \in P^*)$ with the following defining relations:

$$q^0 = 1, q^h q^{h'} = q^{h+h'} \text{ for all } h, h' \in P^*,$$

$$q^h e_i q^{-h} = q^{\langle h,\alpha_i\rangle} e_i, q^h f_i q^{-h} = q^{-\langle h,\alpha_i\rangle} f_i \text{ for all } i \in I, h \in P^*,$$

$$[e_i, f_i] = \delta_{ij} \frac{t_i - t_i^{-1}}{q_i - q_i^{-1}}, \text{ where } q_i = q^{(\alpha_i,\alpha_i)} \text{ and } t_i = q^{(\alpha_i,\alpha_i)h_i} = q_i^{h_i},$$

$$\sum_{k=0}^{b} (-1)^k e_i^{(k)} e_j e_i^{(b-k)} = 0 \text{ for } i \ne j \text{ and } b = 1 - \langle h_i, \alpha_j\rangle.$$

$$\sum_{k=0}^{b} (-1)^k f_i^{(k)} f_j f_i^{(b-k)} = 0 \text{ for } i \ne j \text{ and } b = 1 - \langle h_i, \alpha_j\rangle.$$

Here we use the following notations: $[m]_i = \frac{q_i^m - q_i^{-m}}{q_i - q_i^{-1}}$, $[k]_i! = \prod_{m=1}^{k}[m]_i$, and $e_i^{(k)} = e_i^k/[k]_i!$, $f_i^{(k)} = f_i^k/[k]_i!$. For $k < 0$, we put $e_i^{(k)} = f_i^{(k)} = 0$.

It is well-known that $U_q(\mathfrak{g})$ has a Hopf algebra srtucture with a comultiplication Δ defined by

$$\begin{aligned} \Delta(e_i) &= e_i \otimes t_i^{-1} + 1 \otimes e_i, \\ \Delta(f_i) &= f_i \otimes 1 + t_i \otimes f_i, \\ \Delta(q^h) &= q^h \otimes q^h. \end{aligned}$$

for all $i \in I$, $h \in P^*$. The tensor product of two $U_q(\mathfrak{g})$-modules has a stucture of $U_q(\mathfrak{g})$-module by this comultiplication.

3.1.2 Definition of crystal bases[3]

For $U_q(\mathfrak{g})$-module M and $\lambda \in P$, the λ-weight space of M is defined by $M_\lambda = \{u \in M \mid q^h u = q^{\langle h,\lambda\rangle} u \text{ for all } h \in P^*\}$. For $J \subset I$, let $U_q(\mathfrak{g}_J)$ be the subalgebra of $U_q(\mathfrak{g})$ generated by e_i, f_i, t_i and t_i^{-1} $(i \in J)$. We say that M is *integrable* if $M = \bigoplus_{\lambda\in P} M_\lambda$ and M is a union of finite-dimensional $U_q(\mathfrak{g}_{\{i\}})$-modules for each $i \in I$. By the representation theory of $U_q(\mathfrak{sl}_2)$ any element $u \in M_\lambda$ can be uniquely written as

$$u = \sum_{k\geq 0} f_i^{(k)} u_k,$$

where $u_k \in \ker e_i \cap M_{\lambda+k\alpha_i}$. We define the Kashiwara operators $\tilde{e}_i$ and $\tilde{f}_i$ on M as follows.

$$\tilde{e}_i u = \sum f_i^{(k-1)} u_k,$$

$$\tilde{f}_i u = \sum f_i^{(k+1)} u_k.$$

Let $\mathbf{A}$ be the subring of $\mathbf{Q}(q)$ consisting of the rational functions regular at $q = 0$. A *crystal lattice* of integrable $U_q(\mathfrak{g})$-module M is a free A-submodule of M such that (1) $M \cong \mathbf{Q}(q) \otimes_A L$, (2) $L = \oplus_{\lambda\in P} L_\lambda$ where $L_\lambda = L \cap M_\lambda$, (3) $\tilde{e}_i L \subset L$, $\tilde{f}_i L \subset L$. A *crystal base* of the integrable $U_q(\mathfrak{g})$-module M is a pair (L, B) such that

(i) L is a crystal lattice of M,

(ii) B is a $\mathbf{Q}$-base of L/qL,

(iii) $B = \sqcup_{\lambda\in P} B_\lambda$ where $B_\lambda = B \cap (L_\lambda/qL_\lambda)$,

(iv) $\tilde{e}_i B \subset B \sqcup \{0\}$, $\tilde{f}_i B \subset B \sqcup \{0\}$,

(v) for $b, b' \in B$, $b' \in \tilde{f}_i b$ if and only if $b = \tilde{e}_i b'$ for $i \in I$.

We sometimes replace (ii) by: $B_{ps} = B' \sqcup (-B')$ where B' is a $\mathbf{Q}$-base of L/qL. We call (L, B_{ps}) a crystal psuedo-base and $B_{ps}/\{\pm 1\}$ the associated crystal of (L, B_{ps}).

For a dominant integrable weight $\lambda \in P_+ = \{\lambda \in P \mid \langle h_i, \lambda\rangle \leq 0 \text{ for all } i\}$, let $V(\lambda)$ denote the irreducible integrable $U_q(\mathfrak{g})$-module with highest weight λ. Let u_λ be the highest weight vector of $V(\lambda)$, and let $L(\lambda)$ be the smallest A-submodule of $V(\lambda)$ containing u_λ and stable under $\tilde{f}_i$'s. Set $B(\lambda) = \{b \in L(\lambda)/qL(\lambda) \mid b = \tilde{f}_{i_1} \cdots \tilde{f}_{i_l} u_\lambda \bmod qL(\lambda)\} \setminus \{0\}$. Than $(L(\lambda), B(\lambda))$ is crystal base of $V(\lambda)$. Crystal graph is an oriented colored (by I) graph with $B(\lambda)$ as the set of vertices and colored arrow $b \xrightarrow{i} b'$ if and only if $b' = \tilde{f}_i b$ (hence $\tilde{e}_i b' = b$). Then graph completely describes the actions of $\tilde{e}_i$ and $\tilde{f}_i$ on $B(\lambda)$. For $b \in B(\lambda)$, we set

$$\varepsilon_i(b) = \max\{k \geq 0 \mid \tilde{e}_i^k b \neq 0\},$$

$$\varphi_i(b) = \max\{k \geq 0 \mid \tilde{f}_i^k b \neq 0\}.$$

Note that $B(\lambda) = \sqcup_{\mu\in P} B(\lambda)_\mu$ and for $b \in B(\lambda)_\mu$, we have $\langle h_i, \mathrm{wt}(b)\rangle = \varphi_i(b) - \varepsilon_i(b)$, where $\mathrm{wt}(b) = \mu$ denotes the weight of b.

Proposition 3.1 *Let (L_j, B_j) be crystal bases of integrable $U_q(\mathfrak{g})$-modules M_j $(j = 1, 2)$. Set $L = L_1 \otimes_{\mathbf{A}} L_2 \subset M_1 \otimes M_2$ and $B = \{b_1 \otimes b_2 \mid b_j \in B_j\ (j = 1, 2)\} \subset L/qL\}$. Then (L, B) is the crystal base of $M_1 \otimes M_2$ and the action of $\tilde{f}_i$, $\tilde{e}_i$ is given by*

$$\tilde{f}_i(b_1 \otimes b_2) = \begin{cases} \tilde{f}_i b_1 \otimes b_2 & \text{if } \varphi_i(b_1) > \varepsilon_i(b_2), \\ b_1 \otimes \tilde{f}_i b_2 & \text{if } \varphi_i(b_1) \leq \varepsilon_i(b_2), \end{cases}$$

$$\tilde{e}_i(b_1 \otimes b_2) = \begin{cases} \tilde{e}_i b_1 \otimes b_2 & \text{if } \varphi_i(b_1) \geq \varepsilon_i(b_2), \\ b_1 \otimes \tilde{e}_i b_2 & \text{if } \varphi_i(b_1) < \varepsilon_i(b_2). \end{cases}$$

Corollary 3.2 *For* $b_j \in B_j$ $(j = 1, 2)$, *we have*

$$\begin{aligned}\varepsilon_i(b_1 \otimes b_2) &= \max(\varepsilon_i(b_1), \varepsilon_i(b_2) + \varepsilon_i(b_1) - \varphi_i(b_1)),\\ \varphi_i(b_1 \otimes b_2) &= \max(\varphi_i(b_2), \varphi_i(b_1) + \varphi_i(b_2) - \varepsilon_i(b_2)).\end{aligned}$$

Proposition 3.3 *Let* (L_j, B_j) *be crystal bases of integrable* $U_q(\mathfrak{g})$*-modules* M_j $(j = 1, \ldots, N)$, *and let* $\{a_k\}$ *be the sequence defined by* $a_1 = 0$, $a_{k+1} = a_k + \varphi_i(b_k) - \varepsilon_i(b_{k+1})$. *Then we have*

$$\tilde{f}_i(b_1 \otimes \cdots \otimes b_N) = b_1 \otimes \cdots \otimes b_{k-1} \otimes \tilde{f}_i b_k \otimes b_{k+1} \otimes \cdots \otimes b_N,$$

where $k = max\{j \mid a_j = min\{a_1, \ldots, a_N\}\}$,

$$\tilde{e}_i(b_1 \otimes \cdots \otimes b_N) = b_1 \otimes \cdots \otimes b_{l-1} \otimes \tilde{e}_i b_l \otimes b_{l+1} \otimes \cdots \otimes b_N,$$

where $l = min\{j \mid a_j = min\{a_1, \ldots, a_N\}\}$.

We regard integrable $U_q(\mathfrak{g})$-module M as a union of finite dimensional $U_q(\mathfrak{g}_{\{i\}})$-module for any $i \in I$. Let u_0 be the base of trivial representation of $U_q(\mathfrak{sl}_2)$-module, and let u_+, u_- be a base of 2 dimensional $U_q(\mathfrak{sl}_2)$-module which satisfy

$$fu_+ = u_-,$$

where f is one of generators of $U_q(\mathfrak{sl}_2)$.

In order to calculate the action of $\tilde{e}_i$ and $\tilde{f}_i$ on tensor products, we will use u_+, u_- and u_0. For that purpose we prepare the following. Let B be a crystal base. Let $\mathfrak{W}$ be the set of words generated by u_+, u_-, u_0.

Definition 3.4 *For* $i \in I$ *we define a map*

$$\Psi_i : B \to \mathfrak{W}$$

as follows:

(1) *for* $b \in B$ *such that* $\tilde{e}_i b = \tilde{f}_i b = 0$, *we define*

$$\Psi_i(b) = u_0,$$

(2) *if* b_k *is an element of a maximally connected i-colored crystal subgraph* $b_1 \xrightarrow{i} b_2 \xrightarrow{i} \cdots \xrightarrow{i} b_n$ *where* $n \in \mathbf{Z}_{\geq 2}$, *we define*

$$\Psi_i(b_k) = u_-^{k-1} u_+^{n-k}.$$

We extend Ψ_i *to* $b'_1 \otimes \cdots \otimes b'_N \in B^{\otimes N}$, $N \in \mathbf{Z}_{>0}$, *by*

$$\Psi_i(b'_1 \otimes \cdots \otimes b'_N) = \Psi_i(b'_1) \cdots \Psi_i(b'_N) = u_1 \cdots u_N,$$

where, $\Psi_i(b'_k) = u_k \in \mathfrak{W}$.

For $v = v_1 \cdots v_{N'} \in \mathfrak{W}$, we define the length of the word v by $l(v) = N'$. Let $\mathfrak{N}$ be the set of words generated by $\mathbf{Z}_{>0}$.

Definition 3.5 *We define a map* $\Psi_{\mathfrak{N}} : B^{\otimes N} \to \mathfrak{N}$ *by*

$$\Psi_{\mathfrak{N}}(b'_1 \otimes \cdots \otimes b'_N) = 1^{l(\Psi_i(b_1))} 2^{l(\Psi_i(b_2))} \ldots N^{l(\Psi_i(b_N))}$$

For $r = r_1 \cdots r_{N''} \in \mathfrak{N}$, we also define the length of the word r by $l(r) = N''$.

We put $\mathfrak{W}_j = \{v \in \mathfrak{W} \mid l(v) = j\}$, $\mathfrak{N}_j = \{r \in \mathfrak{N} \mid l(r) = j\}$, where $j \in \mathbf{Z}_{\geq 0}$.

Take words $v = v_1 \cdots v_M \in \mathfrak{W}_M$, $r = r_1 \cdots r_M \in \mathfrak{N}_M$, where $M \in \mathbf{Z}_{\geq 0}$. We put

$$s = \sharp\{s' \mid v_{s'} = u_0, 1 \leq s' \leq M\},$$

$$\{m_1, m_2, \ldots, m_s\} = \{s' \mid v_{s'} = u_0, 1 \leq s' \leq M\},$$

where $1 \leq m_1 < m_2 < \cdots < m_s \leq M$.

Definition 3.6 *We define a map*

$$\mathrm{Red}_0 : \bigcup_{j \in \mathbf{Z}_{\geq 0}} (\mathfrak{W}_j \times \mathfrak{N}_j) \to \bigcup_{j \in \mathbf{Z}_{\geq 0}} (\mathfrak{W}_j \times \mathfrak{N}_j) .$$

The map Red_0 delete all v_{m_i} which equal to u_0 in v and r_{m_i} in r. Namely we can express

$$\mathrm{Red}_0(v,r) = (v_1 \cdots v_{m_1-1} v_{m_1+1} \cdots v_{m_s-1} v_{m_s+1} \cdots v_M, r_1 \cdots r_{m_1-1} r_{m_1+1} \cdots r_{m_s-1} r_{m_s+1} \cdots r_M).$$

We define

$$s_0 = \begin{cases} \min\{m \mid v_m = u_+, v_{m+1} = u_-\} & \text{if there exists } m \text{ such that } v_m = u_+,\ v_{m+1} = u_-, \\ 0 & \text{otherwise.} \end{cases}$$

Definition 3.7 *We define a map*

$$\mathrm{Red}_+ : \bigcup_{j \in \mathbf{Z}_{\geq 0}} (\mathfrak{W}_j \times \mathfrak{N}_j) \to \bigcup_{j \in \mathbf{Z}_{\geq 0}} (\mathfrak{W}_j \times \mathfrak{N}_j) ,$$

for $v = v_1 \cdots v_M \in \mathfrak{W}_M$, $r = r_1 \cdots r_M \in \mathfrak{N}_M$ by

$$\mathrm{Red}_+(v,r) = \begin{cases} (v_1 \cdots v_{s_0-1} v_{s_0+2} \cdots v_M, r_1 \cdots r_{s_0-1} r_{s_0+2} \cdots r_M) & \textit{if } s_0 \geq 1, \\ (v,r) & \textit{if } s_0 = 0. \end{cases}$$

Definition 3.8 *We define a map*

$$\mathrm{Red} : \bigcup_{j \in \mathbf{Z}_{\geq 0}} (\mathfrak{W}_j \times \mathfrak{N}_j) \to \bigcup_{j \in \mathbf{Z}_{\geq 0}} (\mathfrak{W}_j \times \mathfrak{N}_j)$$

by

$$\mathrm{Red} = \mathrm{Red}_+^n \mathrm{Red}_0,$$

where $n \in \mathbf{Z}_{\geq 0}$ such that $n = \min\{n' \mid Red_+^{n'+1}(Red_0(v,r)) = Red_+^{n'}(Red_0(v,r))\}$.

By Definition 3.8, we can denote

$$\mathrm{Red}\,(\Psi_i(b), \Psi_{\mathfrak{N}}(b)) = (v'_1 v'_2 \cdots v'_{M'}, r'_1 r'_2 \cdots r'_{M'})$$

where $M' \in \mathbf{Z}_{\geq 0}$. Let $\mathrm{Pr}_{\mathfrak{W}}$, $\mathrm{Pr}_{\mathfrak{N}}$ be a projection of $\cup_{j \in Z}(\mathfrak{W}_j \times \mathfrak{N}_j)$ to $\mathfrak{W}$, $\mathfrak{N}$ respectively. We define a map

$$\mathrm{Red}_i : B^{\otimes N} \to \mathfrak{W},$$

by

$$\mathrm{Red}_i = \mathrm{Pr}_{\mathfrak{W}} \cdot \mathrm{Red}(\Psi_i, \Psi_{\mathfrak{N}})$$

We define a map

$$\mathrm{Red}_{\mathfrak{N}} : B^{\otimes N} \to \mathfrak{N},$$

by

$$\mathrm{Red}_{\mathfrak{N}} = \mathrm{Pr}_{\mathfrak{N}} \cdot \mathrm{Red}(\Psi_i, \Psi_{\mathfrak{N}})$$

We set

$$s_+ = \begin{cases} \min\{s' \mid v'_{s'} = u_+\} & \text{if there exists } s' \text{ such that } v'_{s'} = u_+, \\ 0 & \text{otherwise,} \end{cases}$$

$$s_- = \begin{cases} \max\{s' \mid v'_{s'} = u_-\} & \text{if there exists } s' \text{ such that } v'_{s'} = u_- \\ 0 & \text{otherwise.} \end{cases}$$

Proposition 3.9 *For $b = b_1 \otimes \cdots \otimes b_N \in B^{\otimes N}$, we have*

$$\tilde{f}_i(b) = \begin{cases} b_1 \otimes \cdots \otimes \tilde{f}_i b_{r_{s_+}} \otimes \cdots \otimes b_N & \text{if } s_+ > 0, \\ 0 & \text{if } s_+ = 0, \end{cases}$$

$$\tilde{e}_i(b) = \begin{cases} b_1 \otimes \cdots \otimes \tilde{e}_i b_{r_{s_-}} \otimes \cdots \otimes b_N & \text{if } s_- > 0, \\ 0 & \text{if } s_- = 0. \end{cases}$$

For $b = b_1 \otimes \cdots \otimes b_N \in B^{\otimes N}$, we denote $\Psi_i(b_k) = u_-^{\varepsilon_i(b_k)} u_+^{\varphi_i(b_k)}$. Then we see that Proposition 3.3 and Proposition 3.9 are equivalent.

Remark 3.10 *For $b = b_1 \otimes \cdots \otimes b_N$, we denote* $\mathrm{Red}_i(b) = u_-^{m_1} u_+^{m_2}$ $(m_1,\, m_2 \in \mathbf{Z}_{\geq 0})$. *Then by Definition 3.4 and Proposition 3.9, we see*

$$\varepsilon_i(b) = m_1,$$

$$\varphi_i(b) = m_2.$$

Example 3.11 *We assume $b = b_1 \otimes b_2 \otimes b_3$, $\Psi_i(b_1) = u_1 = u_- u_+ u_+$, $\Psi_i(b_2) = u_2 = u_0$ and $\Psi_i(b_3) = u_3 = u_- u_+$. We calculate $\tilde{f}_i b$ as follows:*

$$\begin{aligned} \Psi_i(b) &= \Psi_i(b_1)\Psi_i(b_2)\Psi_i(b_3) \\ &= u_1 u_2 u_3 \\ &= u_- u_+ u_+ u_0 u_- u_+, \\ \Psi_{\mathfrak{N}}(b) &= 111233, \\ \mathrm{Red}_0(\Psi_i(b), \Psi_{\mathfrak{N}}(b)) &= (u_- u_+ u_+ u_- u_+, 11133), \\ \mathrm{Red}(\Psi_i(b), \Psi_{\mathfrak{N}}(b)) &= (u_- u_+ u_+, 113), \\ \mathrm{Red}_i(b) &= u_- u_+ u_+. \end{aligned}$$

Then we have $s_+ = 2$, and that s_+-th integer of the word 113 *is* 1. *Therefore,*

$$\tilde{f}_i b = \left(\tilde{f}_i b_1\right) \otimes b_2 \otimes b_3.$$

3.1.3 Crystal bases for $U_q(A_2)$-module[6]

Consider the finite dimensional simple Lie algebra of type A_2 with Cartan matrix $\begin{pmatrix} 2 & -1 \\ -1 & 2 \end{pmatrix}$, Let us denote by $\{\alpha_1, \alpha_2\}$ the set of simple roots and $\{h_1, h_2\}$ the set of simple coroots. Define $\Lambda_i \in \mathfrak{h}^*$ $(i = 1, 2)$ by $\Lambda_i(h_j) = \delta_{ij}$, $(j = 1, 2)$. We put $P = \mathbf{Z}\Lambda_1 \oplus \mathbf{Z}\Lambda_2$, $P^* = \mathbf{Z}h_1 \oplus \mathbf{Z}h_2$. Thus any dominant integral weight $\Lambda \in P_+$ is of the form $\Lambda = m\Lambda_1 + n\Lambda_2$, $(n, m \in \mathbf{Z}_+)$. We define a non-degenerate symmetric bilinear form $(,)$ on $\mathfrak{h}^*$ by $(\alpha_1, \alpha_1) = (\alpha_2, \alpha_2) = 1$, $(\alpha_1, \alpha_2) = (\alpha_2, \alpha_1) = -\frac{1}{2}$, $\langle h_i, \alpha_j \rangle$ is the (i, j)-th element of Cartan matrix and set $t_i = q^{h_i}$ for $i = 1, 2$. Then the corresponding quantized universal enveloping algebra $U_q(A_2)$ is the associative algebra over $\mathbf{Q}(q)$ generated by e_i, f_i, t_i, t_i^{-1} $(i = 1, 2)$ satisfying the relations:

$$t_i t_j = t_j t_i, \qquad t_i t_i^{-1} = t_i^{-1} = t_i = 1,$$

$$t_i e_j t_i^{-1} = q^{\langle h_i, \alpha_j \rangle} e_j, t_i f_j t_i^{-1} = q^{-\langle h_i, \alpha_j \rangle} f_j,$$

$$[e_i, f_j] = \delta_{ij} \frac{t_i - t_i^{-1}}{q - q^{-1}},$$

$$e_i^{(2)} e_j - e_i e_j e_i + e_j e_i^{(2)} = f_i^{(2)} f_j - f_i f_j f_i + f_j f_i^{(2)} = 0 \ (i \neq j).$$

Let V be the 3-dimensional $\mathbf{Q}(q)$-vector space with a basis $\left\{\boxed{1}, \boxed{2}, \boxed{3}\right\}$. We define a $U_q(A_2)$-module structure on V as follows:

$$\begin{array}{lll}
e_1\boxed{1} = 0, & e_1\boxed{2} = \boxed{1}, & e_1\boxed{3} = 0, \\
e_2\boxed{1} = 0, & e_2\boxed{2} = 0, & e_2\boxed{3} = \boxed{2}, \\
f_1\boxed{1} = \boxed{2}, & f_1\boxed{2} = 0, & f_1\boxed{3} = 0, \\
f_2\boxed{1} = 0, & f_2\boxed{2} = \boxed{3}, & f_2\boxed{3} = 0, \\
t_1\boxed{1} = q\,\boxed{1}, & t_1\boxed{2} = q^{-1}\,\boxed{2}, & t_1\boxed{3} = \boxed{3}, \\
t_2\boxed{1} = \boxed{1}, & t_2\boxed{2} = q\,\boxed{2}, & t_2\boxed{3} = q^{-1}\,\boxed{3}.
\end{array}$$

The module V is isomorphic to the integrable highest weight $U_q(A_2)$-module $V(\Lambda_1)$ with highest weight Λ_1 and highest weight vector $\boxed{1}$.

Proposition 3.12 *Put*

$$L(\Lambda_1) = \mathbf{A}\boxed{1} \oplus \mathbf{A}\boxed{2} \oplus \mathbf{A}\boxed{3},$$
$$B(\Lambda_1) = \left\{\boxed{1}, \boxed{2}, \boxed{3}\right\}.$$

Then $(L(\Lambda_1), B(\Lambda_1))$ *is the crystal base of* $V(\Lambda_1)$. *The crystal graph* $B(\Lambda_1)$ *is given by:*

$$\boxed{1} \xrightarrow{1} \boxed{2} \xrightarrow{2} \boxed{3}.$$

Define an ordering on the set $B(\Lambda_1)$ by

$$\boxed{1} \prec \boxed{2} \prec \boxed{3}.$$

We write $\begin{array}{|c|}\hline a \\ \hline b \\ \hline\end{array}$ for $\boxed{a} \otimes \boxed{b}$ $(a \prec b)$. Similar to the $B(\Lambda_1)$-case, the crystal graph $B(\Lambda_2)$ is given by:

$$\begin{array}{|c|}\hline 1 \\ \hline 2 \\ \hline\end{array} \xrightarrow{2} \begin{array}{|c|}\hline 1 \\ \hline 3 \\ \hline\end{array} \xrightarrow{2} \begin{array}{|c|}\hline 2 \\ \hline 3 \\ \hline\end{array}.$$

Proposition 3.13 *The crystal bases of* $B(m\Lambda_1 + n\Lambda_2)$ *is given by the set of semi-standard Young tableaux of the same shape*

$$\begin{aligned} & B(m\Lambda_1 + n\Lambda_2) \\ = & \left\{ \begin{array}{|c|c|c|c|c|c|}\hline b_1^{m+n} & \cdots & b_1^{m+1} & b_1^m & \cdots & b_1^1 \\ \hline b_2^{m+n} & \cdots & b_2^{m+1} & \multicolumn{3}{c}{} \\ \cline{1-3}\end{array} \;\middle|\; b_i^j \in \{1,2,3\}, b_i^j \leq b_i^{j-1}, b_1^j < b_2^j \right\}. \end{aligned}$$

In particular, highest weight element $\hat{b} \in B(m\lambda_1 + n\Lambda_2)$ *is given by*

$$\hat{b} = \begin{array}{|c|c|c|c|c|c|}\hline 1 & \cdots & 1 & 1 & \cdots & 1 \\ \hline 2 & \cdots & 2 & \multicolumn{3}{c}{} \\ \cline{1-3}\end{array}.$$

By Definition 3.4 we have

$$\Psi_i\left(\boxed{j}\right) = \begin{cases} u_+ & \text{if } i = j, \\ u_- & \text{if } i+1 = j, \\ u_0 & \text{otherwise.} \end{cases}$$

An element b of $B(m\Lambda_1 + n\Lambda_2)$ can be expressed in the form

$$\boxed{b_1^1} \otimes \cdots \boxed{b_1^m} \otimes \left(\boxed{b_1^{m+1}} \otimes \boxed{b_2^{m+1}}\right) \otimes \cdots \otimes \left(\boxed{b_1^{m+n}} \otimes \boxed{b_2^{m+n}}\right) \in B(\Lambda_1)^{\otimes m+2n}.$$

Then using Proposition 3.9, we have the action of $\tilde{e}_i$ and $\tilde{f}_i$.

3.1.4 Crystal bases for $U_q(G_2)$-module[7]

Consider the finite dimensional simple Lie algebra of type G_2 with Cartan matrix $\begin{pmatrix} 2 & -1 \\ -3 & 2 \end{pmatrix}$. Let $\{\alpha_1, \alpha_2\}$, $\{h_1, h_2\}$ be the set of simple root and simple coroot respectively. Let $\mathfrak{h}$, Λ_i $(i = 1, 2)$, P and P^* be Caltan subalgebra, fundermental weight, weight lattice and dual lattice, respectively. We define a non-degenerate symmetric bilinear form $(,)$ on $\mathfrak{h}^*$ by $(\alpha_1, \alpha_1) = 3$, $(\alpha_2, \alpha_2) = 1$, $(\alpha_1, \alpha_2) = (\alpha_2, \alpha_1) = -\frac{3}{2}$, $\langle h_i, \alpha_j \rangle$ is (i, j)-element of Cartan matrix and set $q_i = q^{(\alpha_i, \alpha_i)}$, $t_i = q^{h_i}$ for $i = 1, 2$. Then the corresponding quantized universal enveloping algebra $U_q(G_2)$ is the associative algebra over $\mathbf{Q}(q)$ generated by e_i, f_i, t_i, t_i^{-1} $(i = 1, 2)$ satisfying the relations:

$$t_i t_j = t_j t_i, \qquad t_i t_i^{-1} = t_i^{-1} = t_i = 1,$$

$$t_i e_j t_i^{-1} = q_i^{\langle h_i, \alpha_j \rangle} e_j, t_i f_j t_i^{-1} = q_i^{-\langle h_i, \alpha_j \rangle} f_j,$$

$$[e_i, f_j] = \delta_{ij} \frac{t_i - t_i^{-1}}{q_i - q_i^{-1}},$$

$$e_1^{(2)} e_2 - e_1 e_2 e_1 + e_2 e_1^{(2)} = f_1^{(2)} f_2 - f_1 f_2 f_1 + f_2 f_1^{(2)} = 0,$$

$$\sum_{n=0}^{4} (-1)^n e_2^{(n)} e_1 e_2^{(4-n)} = \sum_{n=0}^{4} (-1)^n e_2^{(n)} e_1 e_2^{(4-n)} = 0.$$

Let V be the 14-dimensional $\mathbf{Q}(q)$-vector space with a basis

$$\left\{ \boxed{i}, \boxed{\overline{i}} \;\middle|\; i = 1, \ldots, 6 \right\} \cup \left\{ \boxed{0_1}, \boxed{0_2} \right\}.$$

We define a $U_q(G_2)$-module structure on V as follows:

$e_1 \boxed{2} = \boxed{1}$, $e_1 \boxed{5} = \boxed{4}$, $e_1 \boxed{0_2} = [2]_2 \boxed{6}$, $e_1 \boxed{\overline{6}} = \boxed{0_2} + \frac{1}{[2]_2} \boxed{0_1}$,

$e_1 \boxed{\overline{4}} = \boxed{\overline{5}}$, $e_1 \boxed{\overline{1}} = \boxed{\overline{2}}$,

$e_2 \boxed{3} = [3]_2 \boxed{2}$, $e_2 \boxed{4} = [2]_2 \boxed{3}$, $e_2 \boxed{6} = \boxed{4}$, $e_2 \boxed{0_1} = [2]_2 \boxed{5}$,

$e_2 \boxed{\overline{5}} = \boxed{0_1} + \frac{[3]_2}{[2]_1} \boxed{0_2}$, $e_2 \boxed{\overline{4}} = [3]_2 \boxed{\overline{6}}$, $e_2 \boxed{\overline{3}} = [2]_2 \boxed{\overline{4}}$, $e_2 \boxed{\overline{2}} = \boxed{\overline{3}}$,

$f_1 \boxed{1} = \boxed{2}$, $f_1 \boxed{4} = \boxed{5}$, $f_1 \boxed{6} = \boxed{0_2} + \frac{1}{[2]_2} \boxed{0_1}$,

$f_1 \boxed{0_2} = [2]_2 \boxed{\overline{6}}$, $f_1 \boxed{\overline{5}} = \boxed{\overline{4}}$, $f_1 \boxed{\overline{2}} = \boxed{\overline{1}}$,

$f_2 \boxed{2} = \boxed{3}$, $f_2 \boxed{3} = [2]_2 \boxed{4}$, $f_2 \boxed{4} = [3]_2 \boxed{6}$, $f_2 \boxed{5} = \boxed{0_1} + \frac{[3]_2}{[2]_1} \boxed{0_2}$,

$f_2 \boxed{0_1} = [2]_2 \boxed{\overline{5}}$, $f_2 \boxed{\overline{6}} = \boxed{\overline{4}}$, $f_2 \boxed{\overline{4}} = [2]_2 \boxed{\overline{3}}$, $f_2 \boxed{\overline{3}} = [3]_2 \boxed{\overline{2}}$,

$t_1 \boxed{1} = q_1 \boxed{1}$, $t_1 \boxed{2} = q_1^{-1} \boxed{2}$, $t_1 \boxed{3} = \boxed{3}$, $t_1 \boxed{4} = q_1 \boxed{4}$, $t_1 \boxed{5} = q_1^{-1} \boxed{5}$,

$t_1 \boxed{6} = q_1^{-2} \boxed{6}$, $t_1 \boxed{0_1} = \boxed{0_1}$, $t_1 \boxed{0_2} = \boxed{0_2}$, $t_1 \boxed{\overline{6}} = q_1^{-2} \boxed{\overline{6}}$,

$t_1 \boxed{\overline{5}} = q_1 \boxed{\overline{5}}$, $t_1 \boxed{\overline{4}} = q_1^{-1} \boxed{\overline{4}}$, $t_1 \boxed{\overline{3}} = \boxed{\overline{3}}$, $t_1 \boxed{2} = q_1 \boxed{2}$, $t_1 \boxed{\overline{1}} = q_1^{-1} \boxed{\overline{1}}$,

$t_2 \boxed{1} = \boxed{1}$, $t_2 \boxed{2} = q_2^3 \boxed{2}$, $t_2 \boxed{3} = q_2 \boxed{3}$, $t_2 \boxed{4} = q_2^{-1} \boxed{4}$, $t_2 \boxed{5} = q_2^2 \boxed{5}$,

$t_2 \boxed{6} = q_2^{-3} \boxed{6}$, $t_2 \boxed{0_1} = \boxed{0_1}$, $t_2 \boxed{0_2} = \boxed{0_2}$, $t_2 \boxed{\overline{6}} = q_2^3 \boxed{\overline{6}}$,

$t_2 \boxed{\overline{5}} = q_2^{-2} \boxed{\overline{5}}$, $t_2 \boxed{\overline{4}} = q_2 \boxed{\overline{4}}$, $t_2 \boxed{\overline{3}} = q_2^{-1} \boxed{\overline{3}}$, $t_2 \boxed{\overline{2}} = q_2^{-3} \boxed{\overline{2}}$, $t_2 \boxed{\overline{1}} = \boxed{\overline{1}}$.

The other cases, $b \in B(\Lambda_1)$ is annihilated by $\tilde{e}_i$ or $\tilde{f}_i$.

The module V is isomorphic to the integrable highest weight $U_q(G_2)$-module $V(\Lambda_1)$ with highest weight Λ_1 and highest weight vector $\boxed{1}$.

Proposition 3.14 *Put*

$$L(\Lambda_1) = \bigoplus_{i=1}^{6} \left(\mathbf{A}\boxed{i} \oplus \mathbf{A}\boxed{\overline{i}} \right) \bigoplus \mathbf{A}\boxed{0_1} \bigoplus \mathbf{A}\boxed{0_2},$$

$$B(\Lambda_1) = \left\{ \boxed{i}\,,\ \mathbf{A}\boxed{\overline{i}} \,\middle|\, i = 1, 2, \ldots, 6 \right) \cup \left\{ \boxed{0_1},\ \boxed{0_2} \right\}.$$

Then $(L(\Lambda_1), B(\Lambda_1))$ is the crystal base of $V(\Lambda_1)$. The crystal graph $B(\Lambda_1)$ is given by:



.

Define an ordering on the set $B(\Lambda_1)$ by

$$\boxed{1} \prec \boxed{2} \prec \boxed{3} \prec \boxed{4} \prec \begin{array}{c}\boxed{5}\\ \boxed{6}\end{array} \prec \begin{array}{c}\boxed{0_1}\\ \boxed{0_2}\end{array} \prec \begin{array}{c}\boxed{\overline{5}}\\ \boxed{\overline{6}}\end{array} \prec \boxed{\overline{4}} \prec \boxed{\overline{3}} \prec \boxed{\overline{2}} \prec \boxed{\overline{1}}.$$

We define the map sgn, x_i, and $\overline{x}_i$.

For $a \in \mathbf{R}$, we define $\operatorname{sgn}(a)$ by

$$\operatorname{sgn}(a) = \begin{cases} \dfrac{a}{|a|} & a \neq 0, \\ 0 & a = 0. \end{cases}$$

For $b = \boxed{b_1}\boxed{\cdots}\boxed{b_n}$, we define $x_i(b)$, $\overline{x}_i(b)$ by

$$x_i(b) = \sharp\left\{ \boxed{b_k} = \boxed{i} \,\middle|\, k = 1, \ldots, n \right\} \quad (i = 1, \ldots, 6, 0_1, 0_2),$$

$$\overline{x}_i(b) = \sharp\left\{ \boxed{b_k} = \boxed{\overline{i}} \,\middle|\, k = 1, \ldots, n \right\} \quad (i = 1, \ldots, 6).$$

Proposition 3.15 *The crystal bases of $B(n\Lambda_1)$ is given by following set of restrected semi-standard Young tableaux:*

$$\left\{ \begin{array}{ccc|l} b & = & \boxed{b_1}\boxed{\cdots}\boxed{b_l} & b_k \preceq b_{k+1}\ (k = 1, \cdots, l-1) \\ & & & x_5(b) + x_{0_1}(b) + x_{0_2} + \overline{x}_5(b) \leq 1, \\ & & & x_3(b) + x_4(b) + x_5(b) \leq 1, \\ & = & \boxed{b_l} \otimes \cdots \otimes \boxed{b_1} & \overline{x}_5(b) + \overline{x}_4(b) + \overline{x}_3(b) \leq 1, \\ & & & x_5(b) + \operatorname{sgn}(x_6(b)) + x_{0_1}(b) \leq 1, \\ & & & x_{0_1}(b) + \operatorname{sgn}(\overline{x}_6(b)) + \overline{x}_5(b) \leq 1 \end{array} \right\}.$$

By Definition 3.4, we have

$$\begin{aligned} &\Psi_1\left(\boxed{1}\right) = u_+, \quad \Psi_1\left(\boxed{2}\right) = u_-, \quad \Psi_1\left(\boxed{3}\right) = u_0, \quad \Psi_1\left(\boxed{4}\right) = u_+, \\ &\Psi_1\left(\boxed{5}\right) = u_-, \quad \Psi_1\left(\boxed{6}\right) = u_+^2, \quad \Psi_1\left(\boxed{0_1}\right) = u_0, \quad \Psi_1\left(\boxed{0_2}\right) = u_-u_+, \\ &\Psi_1\left(\boxed{\overline{6}}\right) = u_-^2, \quad \Psi_1\left(\boxed{\overline{5}}\right) = u_+, \quad \Psi_1\left(\boxed{\overline{4}}\right) = u_-, \quad \Psi_1\left(\boxed{\overline{3}}\right) = u_0, \\ &\Psi_1\left(\boxed{\overline{2}}\right) = u_+, \quad \Psi_1\left(\boxed{\overline{1}}\right) = u_-, \end{aligned}$$

$$\begin{aligned}
&\Psi_2\left(\boxed{1}\right) = u_0, \quad \Psi_2\left(\boxed{2}\right) = u_+^3, \quad \Psi_2\left(\boxed{3}\right) = u_- u_+^2, \quad \Psi_2\left(\boxed{4}\right) = u_-^2 u_+,\\
&\Psi_2\left(\boxed{5}\right) = u_+^2, \quad \Psi_2\left(\boxed{6}\right) = u_-^3, \quad \Psi_2\left(\boxed{0_1}\right) = u_- u_+, \quad \Psi_2\left(\boxed{0_2}\right) = u_0,\\
&\Psi_2\left(\boxed{\bar{6}}\right) = u_+^3, \quad \Psi_2\left(\boxed{\bar{5}}\right) = u_-^2, \quad \Psi_2\left(\boxed{\bar{4}}\right) = u_- u_+^2, \quad \Psi_2\left(\boxed{\bar{3}}\right) = u_-^2 u_+,\\
&\Psi_2\left(\boxed{\bar{2}}\right) = u_-^3, \quad \Psi_2\left(\boxed{\bar{1}}\right) = u_0.
\end{aligned} \tag{3.1.1}$$

3.1.5 Affine crystal

Here we recall the definition of affine crystal. Let $\mathfrak{g}$ be an affine Lie algebra over $\mathbf{Q}$ with an indecomposable generalized Cartan Matrix. Let $c \in \sum_i \mathbf{Z}_{\geq 0}\mathfrak{h}_i$ be the canonical central element. Let $\{\alpha_i \mid i \in I\} \subset \mathfrak{h}^*$ be the set of simple roots, and let $\{h_i \mid i \in I\} \subset \mathfrak{h}$ be the set of coroots. Let $\delta \in \sum_i \mathbf{Z}_{\geq 0}\alpha_i$ be the generator of null roots. Set $\mathfrak{h}_{cl} = \bigoplus_{i \in I} \mathbf{Q}h_i \subset \mathfrak{h}$ and $\mathfrak{h}_{cl}^* = (\bigoplus_{i \in I} \mathbf{Q}h_i)^*$. Let $cl : \mathfrak{h}^* \to \mathfrak{h}_{cl}^*$ denote the canonical morphism. We have an exact sequence $0 \to \mathbf{Q}\delta \to \mathfrak{h}^* \to \mathfrak{h}_{cl}^* \to 0$. Fix $i_0 \in I$ and take an integer d such that $\delta - d\alpha_{i_0} \in \sum_{i \neq i_0} \mathbf{Z}\alpha_i$. For simplicity we write 0 for i_0. We define a map $af : \mathfrak{h}_{cl}^* \to \mathfrak{h}^*$ satisfying: $cl \circ af = \mathrm{id}$ and $cl \circ af(\alpha_i) = \alpha_i$ for $i \neq 0$. Let Λ_i be the element of $\mathfrak{h}_{cl}^* \subset \mathfrak{h}^*$ such that $\langle h_j, \Lambda_i \rangle = \delta_{ij}$. Hence we have $\alpha_i = \sum_j \langle h_j, \alpha_i \rangle af(\Lambda_i) + \delta_{i,0} d^{-1}\delta$. We define $P = \sum_i \mathbf{Z}af(\Lambda_i) + \mathbf{Z}d^{-1}\delta \subset \mathfrak{h}^*$ and $P_{cl} = cl(P) \subset \mathfrak{h}_{cl}^*$. An element of P is called an *affine weight* and an element in P_{cl} is called a *classical weight*. Let $U_q'(\mathfrak{g})$ be the quantized universal enveloping algebra associated with P_{cl}. A P-weighted crystal is called an *affine crystal* and a P_{cl}-weighted crystal is called a *classical crystal*. Let $\mathrm{Mod}^f(\mathfrak{g}, P_{cl})$ be the category of $U_q'(\mathfrak{g})$-module M satisfing the following conditions:

$$M \text{ has the weight decomposition } M = \oplus_{\lambda \in P_{cl}} M_\lambda,$$

and

$$M \text{ is finite-dimensional over } \mathbf{Q}(q).$$

For a $U_q'(\mathfrak{g})$-module M in $\mathrm{Mod}^f(\mathfrak{g}, P_{cl})$, we define the $U_q(\mathfrak{g})$-module $\mathrm{Aff}(M)$ by

$$\mathrm{Aff}(M) = \oplus_{\lambda \in P} \mathrm{Aff}(M)_\lambda, \ \mathrm{Aff}(M)_\lambda = M_{cl(\lambda)} \text{ for } \lambda \in P.$$

The actions of e_i and f_i are defined by the commutative diagrams

$$\begin{tikzcd}
\mathrm{Aff}(M)_\lambda \arrow[r, "e_i"] \arrow[d, phantom, "\wr|"] & \mathrm{Aff}(M)_{\lambda+\alpha_i} \arrow[d, phantom, "\wr|"] & \textit{and} & \mathrm{Aff}(M)_\lambda \arrow[r, "f_i"] \arrow[d, phantom, "\wr|"] & \mathrm{Aff}(M)_{\lambda-\alpha_i} \arrow[d, phantom, "\wr|"] \\
M_{cl(\lambda)} \arrow[r, "e_i"] & M_{cl(\lambda+\alpha_i)}, & & M_{cl(\lambda)} \arrow[r, "f_i"] & M_{cl(\lambda-\alpha_i)}.
\end{tikzcd}$$

We define the $U_q'(\mathfrak{g})$-linear automorphism T of $\mathrm{Aff}(M)$ by

$$\begin{tikzcd}
\mathrm{Aff}(M)_\lambda \arrow[r, "T"] \arrow[d, phantom, "\wr|"] & \mathrm{Aff}(M)_{\lambda+\delta} \arrow[d, phantom, "\wr|"] \\
M_{cl(\lambda)} \arrow[r, "id"] & M_{cl(\lambda+\delta)}.
\end{tikzcd}$$

3.1.6 Quantized universal enveloping algebra $U_q(G_2^{(1)})$-module

Consider the finite dimensional affine Lie algebra $G_2^{(1)}$ with Cartan matrix $\begin{pmatrix} 2 & -1 & 0 \\ -1 & 2 & -1 \\ 0 & -3 & 2 \end{pmatrix}$. Let $\{\alpha_0, \alpha_1, \alpha_2\}$, $\{h_0, h_1, h_2\}$ be the set of simple root and simple coroot respectively. Let $\mathfrak{h}$ and Λ_i $(i = 0, 1, 2)$ be Caltan subalgebra, fundermental weight, respectively. We define a non-degenerate symmetric bilinear form $(,)$ on $\mathfrak{h}^*$ by $(\alpha_0, \alpha_0) = (\alpha_1, \alpha_1) = 3$, $(\alpha_2, \alpha_2) = 1$, $(\alpha_0, \alpha_1) = (\alpha_1, \alpha_0) = (\alpha_1, \alpha_2) = (\alpha_2, \alpha_1) = -\frac{3}{2}$, $\langle h_i, \alpha_j \rangle$ is (i, j)-element of Cartan matrix and set $q_i = q^{(\alpha_i, \alpha_i)}$, $t_i = q^{h_i}$ for $i = 1, 2$. Canonical central element is $c = h_0 + 2h_1 + h_2$. Then the corresponding quantized universal enveloping algebra $U_q(G_2^{(1)})$ is the associative algebra over $\mathbf{Q}(q)$ generated by e_i, f_i, t_i, t_i^{-1} $(i = 0, 1, 2)$ satisfying the relations:

$$t_i t_j = t_j t_i, \quad t_i t_i^{-1} = t_i^{-1} = t_i = 1,$$

$$t_i e_j t_i^{-1} = q_i^{\langle h_i, \alpha_j \rangle} e_j, t_i f_j t_i^{-1} = q_i^{-\langle h_i, \alpha_j \rangle} f_j,$$

$$[e_i, f_j] = \delta_{ij} \frac{t_i - t_i^{-1}}{q_i - q_i^{-1}}$$

$$\sum_{n=0}^{l} (-1)^n e_i^{(n)} e_j e_i^{(l-n)} = \sum_{n=0}^{l} (-1)^n e_i^{(n)} e_j e_i^{(l-n)} = 0, \quad (i \neq j, \ l = 1 - \langle h_i, \alpha_j \rangle) .$$

3.2 Perfect crystal[9]

Let B be a classical crystal. For $b \in B$, we set $\varepsilon(b) = \sum_i \varepsilon_i(b)\Lambda_i$, and $\varphi(b) = \sum_i \varphi_i(b)\Lambda_i$. Note that $\mathrm{wt}(b) = \varphi(b) - \varepsilon(b)$. Set $P_{cl}^+ = \{\lambda \in P_{cl} \mid \langle h_i, \lambda \rangle \geq 0 \text{ for all } i \in I\}$ and for $l \in \mathbf{Z}_{\geq 0}$, let $(P_{cl}^+)_l = \{\lambda \in P_{cl}^+ \mid \langle c, \lambda \rangle = l\}$.

Definition 3.16 (Definition 4.6.1 in [9])
For $l \in \mathbf{Z}_{\geq 0}$, we say that B is a *perfect crystal of level* l if

(1) $B \otimes B$ is connected.

(2) There exists $\lambda_0 \in P_{cl}$ such that $\mathrm{wt}(B) \in \lambda_0 + \sum_{i \neq 0} \mathbf{Z}_{\leq 0} cl(\alpha_i)$ and that $\sharp(B_{\lambda_0}) = 1$.

(3) There is a finite dimensional $U_q'(\mathfrak{g})$-module with a crystal pseudo-base (L, B_{ps}) such that $B \cong B_{ps}/\pm 1$.

(4) For any $b \in B$, we have $\langle c, \varepsilon(b) \rangle \geq l$.

(5) The maps $\varepsilon, \varphi \ : \ B_l = \{b \in B \mid \langle c, \varepsilon(b) \rangle = l\} \to (P_{cl}^+)_l$ are bijective.

The elements in B_l are called *minimal elements*.

Definition 3.17 *Let M be a $U_q(\mathfrak{g})$-module. A symmetric bilinear form $(,)$ on M is called a prepolarization of M if it satisfies for $u, v \in M$:*

(i) $(q^h u, v) = (u, q^h v)$,

(ii) $(e_i u, v) = (u, q_i^{-1} t_i^{-1} f_i v)$,

(iii) $(f_i u, v) = (u, q_i^{-1} t_i e_i v)$.

A prepolarization is called a polarization if it is positive definite.

Lemma 3.18 (Lemma 3.4.4 in [9])
Let M be a $U_q'(\mathfrak{g})$-module in $Mod^f(\mathfrak{g}, P_{cl})$. Assume that M has a crystal pseudo-base (L, B) such that

(i) *there exists $\lambda \in P_{cl}$ such that $\sharp(B/\{\pm 1\})_\lambda = 1$,*

(ii) *$B/\{\pm 1\}$ is connected.*

Then M is an irreducible $U_q'(\mathfrak{g})$-module.

Let us introduce the subalgebras $\mathbf{A}_Z$ and $\mathbf{K}_Z$ of $\mathbf{Q}(q)$ as follows:

$$\mathbf{A}_Z = \{f(q)/g(q) \mid f(q), \, g(q) \in \mathbf{Z}[q], \, g(0) = 1\},$$

$$\mathbf{K}_Z = \mathbf{A}_Z[q^{-1}].$$

Proposition 3.19 (Proposition 2.6.2 in [8])
Assume that $\mathfrak{g}$ *is finite dimensional and let* M *be a finite dimensional integrable* $U_q(\mathfrak{g})$*-module of* M*. Let* $(,)$ *be a prepolarization on* M*, and* M_{K_Z} *a* $U_q(\mathfrak{g})$*-submodule of* M *such that* $(M_{K_Z}, M_{K_Z}) \subset K_Z$*. Let* $\lambda_1, \ldots, \lambda_m \in P_+$ $(\lambda_k = \sum_j a_{kj}\Lambda_j,\ a_{kj} \in \mathbf{Z}_{\geq 0},\ k = 1, \ldots, m)$*, and we assume the following conditions.*

(i) $\dim M_{\lambda_k} \leq \sum_{j=1}^{m} \dim V(\lambda_j)_{\lambda_k}$ *for* $k = 1, \ldots, m$*.*

(ii) *There exists* $u_j \in (M_{K_Z})_{\lambda_j}$ $(j = 1, \ldots, m)$ *such that* $(u_j, u_k) \in \delta_{jk} + qA$ *and* $(e_i u_j, e_i u_j) \in qq^{-2(1+\langle h_i, \lambda_j\rangle)}A$ *for all* $i \in I$*.*

Set $L = \{U \in M \mid (u, u) \in A\}$ *and set* $B = \{b \in M_{K_Z} \cap L / M_{K_Z} \cap qL \,|(b, b)_0 = 1\}$*. Then we have*

(i) $(,)$ *is a polarization on* M*,*

(ii) $M \cong \oplus V(\lambda_j)$*,*

(iii) $(,)_0$ *is positive definite, and* (L, B) *is a crystal pseudo-base of* M*.*

Lemma 3.20 (Lemma 4.1.2 [8]) *A polarization* $(,)$ *for Kac-Moody Lie algebra with symmetrizable Cartan matrix satisfies following relations.*

(i) *Let* b *be a global base which satisfies* $e_i b = 0$ *and* $\langle h_i, \mathrm{wt}(b)\rangle = 1$*. Then* $(b, b) = (f_i b, f_i b)$*.*

(ii) *Let* b *be a global base which satisfies* $e_i b = 0$ *and* $\langle h_i, \mathrm{wt}(b)\rangle = 2$*. Then* $(b, b) = (f_i^{(2)} b, f_i^{(2)} b) = q_i^{-1}[2]_i^{-1}(f_i b, f_i b)$*.*

(iii) *Let* b *be a global base which satisfies* $e_i b = 0$ *and* $\langle h_i, \mathrm{wt}(b)\rangle = 3$*. Then* $(b, b) = (f_i^{(3)} b, f_i^{(3)} b) = q_i^{-2}[3]_i^{-1}(f_i^{(2)} b, f_i^{(2)}) = q_i^{-1}[2]_i^{-1}(f_i b, f_i b)$*.*

The existace of polarization for Kac-Moody Lie algebra with symmetrizable Cartan matrix is proved in Proposition 3.4.4 [4].

Proposition 3.21 (Proposition 3.4.5 in [8])
Let m *be a positive integer and assume the following conditions:*

(i) $\langle h_i, l\lambda_0 + j\alpha_{i_0}\rangle \geq 0$ *for* $i \neq i_0$ *and* $0 \leq j \leq m$*,*

(ii) $\dim(V_l)_{l\lambda_0 + k\alpha_{i_0}} \leq \sum_{j=0}^{m} \dim V(l\lambda_0 + j\alpha_{i_0})_{l\lambda_0 + k\alpha_{i_0}}$ *for* $0 \leq k \leq m$*, where* $V(\lambda)$ *is an irreducible* $U_q(\mathfrak{g}_{I \setminus i_0})$*-module with highest weight* λ*,*

(iii) *There exists* $i_1 \in I$ *such that* $\{i \in I \mid \langle h_{i_0}, \alpha_i\rangle < 0\} = \{i_1\}$*,*

(iv) $-\langle h_{i_0}, l\lambda_0 - \alpha_{i_1}\rangle \geq 0$*.*

Then we have

$$V_l \cong \oplus_{j=0}^{m} V(l\lambda_0 + j\alpha_{i_0}) \text{ as a } U_q(\mathfrak{g}_{I \setminus \{i_0\}})\text{-module,}$$

and V_l *admits a crystal pseudobase as a* $U_q'(\mathfrak{g})$*-module.*

4 Level-one representation and fusion construction

4.1 Construction of the polarization of V^1

The space $V^1 \cong V(\Lambda_1) \oplus V(0)$ is endowed with a $U_q'(G_2^{(1)})$-module structure as follows.

$f_0 \boxed{6} = \boxed{2}$, $f_0 \boxed{4} = \boxed{3}$, $f_0 \boxed{3} = \boxed{4}$, $f_0 \boxed{2} = \boxed{6}$,

$f_0\boxed{\overline{1}} = \boxed{\cdot} + \frac{1}{[2]_1}\boxed{0_2}$, $f_0\boxed{\cdot} = [2]_0\boxed{1}$,

$f_1\boxed{1} = \boxed{2}$, $f_1\boxed{4} = \boxed{5}$, $f_1\boxed{6} = \boxed{0_2} + \frac{1}{[2]_2}\boxed{0_1} + \frac{1}{[2]_1}\boxed{\cdot}$,

$f_1\boxed{0_2} = [2]_2\boxed{\overline{6}}$, $f_1\boxed{\overline{5}} = \boxed{\overline{4}}$, $f_1\boxed{\overline{2}} = \boxed{\overline{1}}$,

$f_2\boxed{2} = \boxed{3}$, $f_2\boxed{3} = [2]_2\boxed{4}$, $f_2\boxed{4} = [3]_2\boxed{6}$, $f_2\boxed{5} = \boxed{0_1} + \frac{[3]_2}{[2]_1}\boxed{0_2}$,

$f_2\boxed{0_1} = [2]_2\boxed{\overline{5}}$,$f_2\boxed{\overline{6}} = \boxed{\overline{4}}$, $f_2\boxed{\overline{4}} = [2]_2\boxed{\overline{3}}$, $f_2\boxed{\overline{3}} = [3]_2\boxed{\overline{2}}$,

$e_0\boxed{6} = \boxed{\overline{2}}$, $e_0\boxed{4} = \boxed{\overline{3}}$, $e_0\boxed{3} = \boxed{\overline{4}}$, $e_0\boxed{2} = \boxed{\overline{6}}$,

$e_0\boxed{1} = \boxed{\cdot} + \frac{1}{[2]_1}\boxed{0_2}$, $e_0\boxed{\cdot} = [2]_0\boxed{\overline{1}}$,

$e_1\boxed{\overline{1}} = \boxed{\overline{2}}$, $e_1\boxed{\overline{4}} = \boxed{\overline{5}}$, $e_1\boxed{\overline{6}} = \boxed{0_2} + \frac{1}{[2]_2}\boxed{0_1} + \frac{1}{[2]_1}\boxed{\cdot}$,

$e_1\boxed{0_2} = [2]_2\boxed{6}$, $e_1\boxed{5} = \boxed{4}$, $e_1\boxed{2} = \boxed{1}$

$e_2\boxed{\overline{2}} = \boxed{\overline{3}}$, $e_2\boxed{\overline{3}} = [2]_2\boxed{\overline{4}}$, $e_2\boxed{\overline{4}} = [3]_2\boxed{\overline{6}}$, $e_2\boxed{\overline{5}} = \boxed{0_1} + \frac{[3]_2}{[2]_1}\boxed{0_2}$,

$e_2\boxed{0_1} = [2]_2\boxed{5}$, $e_2\boxed{6} = \boxed{4}$, $e_2\boxed{4} = [2]_2\boxed{3}$, $e_2\boxed{3} = [3]_2\boxed{2}$,

$q^{h_0} = q^{-2h_1-h_2}$,

where $\boxed{\cdot}$ is the base of $V(0)$.

We put

$$B^1 = \{\boxed{i}, \boxed{0_1}, \boxed{0_2}, \boxed{\overline{i}}, \boxed{\cdot} \mid i = 1, \dots 6\}.$$

Let $(,)_1$ be the polarization on the $U_q(G_2)$-module $V(\Lambda_1)$. We shall define a symmetric bilinear form $(,)$ on V^1 by

$$\left(\boxed{\cdot}, u\right) = \left(u, \boxed{\cdot}\right) = 0 \quad u \in V(\Lambda_1),$$

$$\left(\boxed{\cdot}, \boxed{\cdot}\right) = q_0[2]_0\left(\boxed{1}, \boxed{1}\right)_1,$$

$$(u, v) = (u, v)_1 \quad u, v \in V(\Lambda_1).$$

By Lemma 3.20, we have

$$\left(\boxed{a_1}, \boxed{a_1}\right) = q_2^{-2}[3]_2^{-1}\left(\boxed{a_2}, \boxed{a_2}\right) = q_2^{-3}[3]_2^{-1}[2]_2^{-1}\left(\boxed{0_1}, \boxed{0_1}\right) = q_1^{-1}[2]_1^{-1}\left(\boxed{0_2}, \boxed{0_2}\right),$$

where $a_1 = 1, 2, 6, \overline{6}, \overline{2}, \overline{1}$, $a_2 = 3, 4, 5, \overline{5}, \overline{4}, \overline{3}$. It follows that $(,)$ is a polarization on the $U'_q(G_2^{(1)})$-module V^1.

Therfore we see that B^1 is a crystal base and Kashiwara operators act as follows.

$\tilde{f}_0\boxed{\overline{6}} = \boxed{2}$, $\tilde{f}_0\boxed{\overline{4}} = \boxed{3}$, $\tilde{f}_0\boxed{\overline{3}} = \boxed{4}$, $\tilde{f}_0\boxed{\overline{2}} = \boxed{6}$, $\tilde{f}_0\boxed{\overline{1}} = \boxed{\cdot}$, $\tilde{f}_0\boxed{\cdot} = [2]_0\boxed{1}$,

$\tilde{f}_1\boxed{1} = \boxed{2}$, $\tilde{f}_1\boxed{4} = \boxed{5}$, $\tilde{f}_1\boxed{6} = \boxed{0_2}$ $\tilde{f}_1\boxed{0_2} = [2]_2\boxed{\overline{6}}$, $\tilde{f}_1\boxed{\overline{5}} = \boxed{\overline{4}}$, $\tilde{f}_1\boxed{\overline{2}} = \boxed{\overline{1}}$,

$\tilde{f}_2\boxed{2} = \boxed{3}$, $\tilde{f}_2\boxed{3} = [2]_2\boxed{4}$, $\tilde{f}_2\boxed{4} = [3]_2\boxed{6}$, $\tilde{f}_2\boxed{5} = \boxed{0_1}$, $\tilde{f}_2\boxed{0_1} = [2]_2\boxed{\overline{5}}$, $\tilde{f}_2\boxed{\overline{6}} = \boxed{\overline{4}}$,

$\tilde{f}_2\boxed{\overline{4}} = [2]_2\boxed{\overline{3}}$, $\tilde{f}_2\boxed{\overline{3}} = [3]_2\boxed{\overline{2}}$,

$\tilde{e}_0\boxed{6} = \boxed{\overline{2}}$, $\tilde{e}_0\boxed{4} = \boxed{\overline{3}}$, $\tilde{e}_0\boxed{3} = \boxed{\overline{4}}$, $\tilde{e}_0\boxed{2} = \boxed{\overline{6}}$, $\tilde{e}_0\boxed{1} = \boxed{\cdot}$, $\tilde{e}_0\boxed{\cdot} = [2]_2\boxed{\overline{1}}$,

$\tilde{e}_1\boxed{\overline{1}} = \boxed{\overline{2}}$, $\tilde{e}_1\boxed{\overline{4}} = \boxed{\overline{5}}$, $\tilde{e}_1\boxed{\overline{6}} = \boxed{0_2}$, $\tilde{e}_1\boxed{0_2} = [2]_2\boxed{6}$, $\tilde{e}_1\boxed{5} = \boxed{4}$, $\tilde{e}_1\boxed{2} = \boxed{1}$

$\tilde{e}_2\boxed{\overline{2}} = \boxed{\overline{3}}$, $\tilde{e}_2\boxed{\overline{3}} = [2]_2\boxed{\overline{4}}$, $\tilde{e}_2\boxed{\overline{4}} = [3]_2\boxed{\overline{6}}$, $\tilde{e}_2\boxed{\overline{5}} = \boxed{0_1}$, $\tilde{e}_2\boxed{0_1} = [2]_2\boxed{5}$, $\tilde{e}_2\boxed{6} = \boxed{4}$,

$\tilde{e}_2\boxed{4} = [2]_2\boxed{3}$, $\tilde{e}_2\boxed{3} = [3]_2\boxed{2}$,

By checking of conditions of Definition 3.16, we have following proposition.

Proposition 4.1 *B^1 is a perfect crystal of level* 1.

4.2 Decomposition of the tensor product

The tensor product $V^1 \otimes V^1$ is decomposed in the following way:

$$\begin{aligned}&(V(\Lambda_1) \oplus V(0)) \otimes (V(\Lambda_1) \oplus V(0)) \\ &\qquad \cong V(2\Lambda_1) \oplus V(3\Lambda_2) \oplus V(2\Lambda_2) \oplus V(\Lambda_1)^{\oplus 3} \oplus V(0)^{\oplus 2}.\end{aligned} \tag{4.2.1}$$

Lemma 4.2 Following vectors $u_{2\Lambda_1}$, $u_{3\Lambda_2}$, $u_{2\Lambda_2}$, $u^1_{\Lambda_1}$, $u^2_{\Lambda_1}$, $u^3_{\Lambda_1}$, u^1_0 and u^2_0 are the highest weight vectors with the weights $2\Lambda_1$, $3\Lambda_2$, $2\Lambda_2$, Λ_1, Λ_1, Λ_1, 0 and 0 respectively.

$$\begin{aligned}
u_{2\Lambda_1} &= \boxed{1} \otimes \boxed{1}, \\
u_{3\Lambda_2} &= \boxed{1} \otimes \boxed{2} - q^3 \boxed{2} \otimes \boxed{1}, \\
u_{2\Lambda_2} &= \boxed{1} \otimes \boxed{5} - q^3 \boxed{2} \otimes \boxed{4} + \frac{[2]_2}{[3]_2} q^4 \boxed{3} \otimes \boxed{3} - q^7 \boxed{4} \otimes \boxed{2} + q^{10} \boxed{5} \otimes \boxed{1}, \\
u^1_{\Lambda_1} &= \boxed{1} \otimes \boxed{\cdot}, \\
u^2_{\Lambda_1} &= \boxed{\cdot} \otimes \boxed{1}, \\
u^3_{\Lambda_1} &= \boxed{1} \otimes \boxed{0_2} - [2]_1 q^6 \boxed{2} \otimes \boxed{6} + \frac{[2]_1}{[3]_2} q^5 \boxed{3} \otimes \boxed{4} - \frac{[2]_1}{[3]_2} q^6 \boxed{4} \otimes \boxed{3} \\
&\quad + [2]_1 q^9 \boxed{6} \otimes \boxed{2} - q^{12} \boxed{0_2} \otimes \boxed{1}, \\
u^1_0 &= \boxed{\cdot} \otimes \boxed{\cdot}, \\
u^2_0 &= \boxed{1} \otimes \boxed{\overline{1}} - q^3 \boxed{2} \otimes \boxed{\overline{2}} + \frac{1}{[3]_2} q^2 \boxed{3} \otimes \boxed{\overline{3}} - \frac{1}{[3]_2} q^3 \boxed{4} \otimes \boxed{\overline{4}} \\
&\quad + \frac{1}{[3]_2} q^6 \boxed{5} \otimes \boxed{\overline{5}} + q^6 \boxed{6} \otimes \boxed{\overline{6}} - \frac{q^6}{[2]_2[3]_2} \boxed{0_1} \otimes \boxed{0_1} - \frac{q^6}{[2]_1} \boxed{0_2} \otimes \boxed{0_2} \\
&\quad + \frac{q^8}{[3]_2} \boxed{\overline{5}} \otimes \boxed{5} + q^{12} \boxed{\overline{6}} \otimes \boxed{6} - \frac{q^{11}}{[3]_2} \boxed{\overline{4}} \otimes \boxed{4} + \frac{q^{12}}{[3]_2} \boxed{\overline{3}} \otimes \boxed{3} \\
&\quad - q^{15} \boxed{\overline{2}} \otimes \boxed{2} + q^{18} \boxed{\overline{1}} \otimes \boxed{1} - \frac{q^6}{[2]_1[2]_3} \boxed{0_1} \otimes \boxed{0_2} - \frac{q^6}{[2]_1[2]_3} \boxed{0_2} \otimes \boxed{0_1} \\
&\quad - \frac{q^6}{[2]_1^2} \boxed{0_2} \otimes \boxed{\cdot} - \frac{q^6}{[2]_1^2} \boxed{\cdot} \otimes \boxed{0_2}.
\end{aligned}$$

4.3 Calculation of the R-matrix

Let V^1_x be $U'_q(\mathfrak{g})$-module $\mathbf{Q}[x, x^{-1}] \otimes V^1$ with the actions of e_i, f_i, and t_i given by $x^{\delta_{i0}} e_i$, $x^{-\delta_{i0}} f_i$, and t_i, respectively. The R-matrix for V^1 is an invertible $\mathbf{Q}[x, x^{-1}, y, y^{-1}] \otimes U_q(G_2)$-linear map

$$R(x, y) : V_x \otimes V_y \to V_y \otimes V_x$$

satisfying following properties.

(1) $R(x, y) \in \mathbf{Q}(q)[x/y, y/x] \otimes \mathrm{End}_{Q(q)}(V^1 \otimes V^1)$.

(2) $(R(y, z) \otimes 1)(1 \otimes R(x, z))(R(x, y) \otimes 1) = (1 \otimes R(x, y))(R(x, z) \otimes 1)(1 \otimes R(y, z))$.

(3) $R(x, y)R(y, x) \in \mathbf{Q}(q)[x/y, y/x]$.

By the decomposition 4.2.1, $U_q(G_2)$-linearlity and $R = R(x, y)$, we have

$$\begin{aligned}&R(u_{2\Lambda_1}) = a^{2\Lambda_1} u_{2\Lambda_1}, R(u_{3\Lambda_2}) = a^{3\Lambda_2} u_{3\Lambda_2}, R(u_{2\Lambda_2}) = a^{2\Lambda_2} u_{2\Lambda_2}, \\ &R(u^i_{\Lambda_1}) = \sum_{j=1}^{3} a^{\Lambda_1}_{ij} u^j_{\Lambda_1} \ (i = 1, 2, 3), R(u^i_0) = \sum_{j=1}^{2} a^0_{ij} u^j_0 \ (i = 1, 2).\end{aligned} \tag{4.3.1}$$

Consider the highest weight vectors given in Lemma 4.2. Then we have in $V^1_x \otimes V^1_y$:

1. $f_0^{(2)}f_1f_2^{(3)}f_1u^1_{\Lambda_1}=[2]_1q^{-3}x^{-1}y^{-1}u_{2\Lambda_1}$,
2. $f_0^{(2)}f_1f_2^{(3)}f_1u^2_{\Lambda_1}=[2]_1q^{-3}x^{-1}y^{-1}u_{2\Lambda_1}$,
3. $f_0^{(2)}f_1f_2^{(3)}f_1u^3_{\Lambda_1}=[2]_1(x-q^6y)(x+q^{12}y)q^{-6}x^{-2}y^{-2}u_{2\Lambda_1}$,
4. $e_1e_2^{(3)}e_1^{(2)}e_2^{(3)}e_0e^{(2)}e_2^{(3)}e_1e_0u^1_{\Lambda_1}=[2]_1y(x+y)u_{2\Lambda_1}$,
5. $e_1e_2^{(3)}e_1^{(2)}e_2^{(3)}e_0e^{(2)}e_2^{(3)}e_1e_0u^2_{\Lambda_1}=[2]_1q^{-6}x(x+y)u_{2\Lambda_1}$,
6. $e_1e_2^{(3)}e_1^{(2)}e_2^{(3)}e_0e^{(2)}e_2^{(3)}e_1e_0u^3_{\Lambda_1}=[2]_2q^5(q^4+1)(x-q^6y)u_{2\Lambda_1}$,
7. $f_0u^1_{\Lambda_1}=[2]_1q^{-6}y^{-1}u_{2\Lambda_1}$,
8. $f_0u^2_{\Lambda_1}=[2]_1x^{-1}u_{2\Lambda_1}$,
9. $f_0u^3_{\Lambda_1}=0$,
10. $f_0^{(2)}u^1_0=[2]_1^2q^{-3}x^{-1}y^{-1}u_{2\Lambda_1}$,
11. $f_0^{(2)}u^2_0=q^{-12}(x^2+q^{30}y^2)x^{-2}y^{-2}u_{2\Lambda_1}$,
12. $f_1^{(2)}f_2^{(6)}f_1^{(4)}f_2^{(6)}f_1^{(2)}f_0^{(2)}f_1f_2^{(2)}f_1f_2f_0f_1f_2^{(3)}f_1f_0u^1_0$
    $=[2]_1^2(q^4+q^2+1)q^{-8}(q^6y^2+x^2)xyu_{2\Lambda_1}$,
13. $f_1^{(2)}f_2^{(6)}f_1^{(4)}f_2^{(6)}f_1^{(2)}f_0^{(2)}f_1f_2^{(2)}f_1f_2f_0f_1f_2^{(3)}f_1f_0u^2_0=$
    $[2]_2q^{-10}(q^4+q^2+1)((q^{22}+q^{18})y^2+(q^6+1)(q^{16}-q^{14}+q^{12}-2q^{10}+q^8-2q^6+q^4-q^2+1)xy+$
    $(q^4+1)x^2)xyu_{\Lambda_1}$,
14. $f_0^{(2)}f_1f_2^{(3)}f_1^{(2)}f_2^{(3)}u_{3\Lambda_2}=q^{-3}(x-q^6y)(x+y)x^{-2}y^{-2}u_{2\Lambda_1}$,
15. $f_0^{(2)}f_1f_2^{(3)}f^1f_2^{(2)}u_{2\Lambda_2}=q^{-8}(q^4+q^2+1)(x-q^6y)(x-q^{10}y)x^{-2}y^{-2}u_{2\Lambda_1}$.

From these equations and the relation $[R,\Delta(X)]=0$ $(X\in U_q(G_2))$, we obtain
$a_{2\Lambda_1}(q^6y,\ x,\ (q^4+1)[2]_2q^4(x-q^6y))=(q^6x,\ y,\ (q^4+1)[2]_2q^4(y-q^6x))\,{}^t(a^{\Lambda_1}_{ij})$,

$a_{2\Lambda_1}(x,\ y,\ 0)=(y,\ x,\ 0)\,{}^t(a^{\Lambda_1}_{ij})$,

$a_{2\Lambda_1}([2]_1^2q^6,\ (x^2+q^{30}y^2)q^{-3}x^{-1}y^{-1})=([2]_1^2q^6,\ (y^2+q^{30}x^2)q^{-3}x^{-1}y^{-1})\,{}^t(a^0_{ij})$,

$a_{2\Lambda_1}(s_1(x,y),\ s_2(x,y))=(s_1(y,x),\ s_2(y,x))\,{}^t(a^0_{ij})$,

$(x-q^6y)a_{2\Lambda_1}=a_{3\Lambda_2}(y-q^6x)$,

$(x-q^6y)(x-q^{10}y)a_{2\Lambda_1}=a_{2\Lambda_2}(y-q^6x)(y-q^{10}x)$,
where
$s_1(x,y)=[2]_1(q^4-a^2+1)(x^2+q^6y^2)$

$s_2(x,y)=(q^{22}+q^{18})y^2+(q^6+1)(q^{16}-q^{14}+q^{12}-2q^{10}+q^8-2q^6+q^4-q^2+1)xy+(q^4+1)x^2$.
Let $P_{2\Lambda_1}$, $P_{3\Lambda_2}$, $P_{2\Lambda_2}$, $P_{u^i_{\Lambda_1}}$ $(i=1,2,3)$ and $P_{u^i_0}$ $(i=1,2)$ be the projection from $V^1\otimes V^1$ to $V(2\Lambda_1)$, $V(3\Lambda_2)$, $V(2\Lambda_2)$, $U_q(G_2)u^i_{\Lambda_1}$ $(i=1,2,3)$ and $U_q(G_2)u^i_0$ $(i=1,2)$ respectively.

Proposition 4.3 Put $z=xy^{-1}$. Since the R-matrix $R(x,y)$ depends only on x/y, we denote $R(z)=R(x,y)$. Then we have

$$\begin{aligned}
R(z)(u_{2\Lambda_1}) &= (1-q^{12}z)(1-q^{10}z)(1-q^8z)(1-q^6z)u_{2\Lambda_1},\\
R(z)(u_{3\Lambda_2}) &= (1-q^{12}z)(1-q^{10}z)(1-q^8z)(z-q^6)u_{3\Lambda_2},\\
R(z)(u_{2\Lambda_2}) &= (1-q^{12}z)(z-q^{10})(1-q^8z)(z-q^6)u_{2\Lambda_2},
\end{aligned}$$

$$\begin{aligned} R(u^i_{\Lambda_1}) &= \sum_{j=1}^{3} a^{\Lambda_1}_{ij} u^j_{\Lambda_1} \ \ (i=1,2,3), \\ R(u^i_0) &= \sum_{j=1}^{2} a^0_{ij} u^j_0 \ \ (i=1,2). \end{aligned}$$

Here $(a^{\Lambda_1}_{ij})$ and (a^0_{ij}) are given by

$a^{\Lambda_1}_{11} = (1-q^{12}z)(q^6-1)(q^2+1)z(-(q^{16}-q^{14}+q^{12}-q^{10}-q^6)z-(q^4-q^2+1))$,
$a^{\Lambda_1}_{12} = (1-q^{12}z)q^6(1-z)(q^{12}z^2+(q^{12}-q^6-q^4-q^2)z+1)$,
$a^{\Lambda_1}_{13} = (1-q^{12}z)q^3(q^6-1)z(z-1)$,
$a^{\Lambda_1}_{21} = (1-q^{12}z)q^6(1-z)(q^{12}z^2-(q^{10}+q^8+q^6-1)z+1)$,
$a^{\Lambda_1}_{22} = (1-q^{12}z)(q^6-1)(q^2+1)z((-q^{16}+q^{14}-q^{12})z+(q^{10}+q^6-q^4+q^2-1))$,
$a^{\Lambda_1}_{23} = (1-q^{12}z)q^3(q^6-1)(z-1)z$,
$a^{\Lambda_1}_{31} = (1-q^{12}z)q^9(q^{12}-1)(q^4+1)(q^2+1)z(1-z)(z-q^6)$,
$a^{\Lambda_1}_{32} = (1-q^{12}z)q^3(q^{12}-1)(q^4+1)(q^2+1)(1-z)(q^6-z)$,
$a^{\Lambda_1}_{33} = (1-q^{12}z)(z-q^6)(q^6z^2+(q^{18}-q^{12}-q^{10}-q^8-q^6+1)z+q^{12})$,
$a^0_{11} = q^{30}z^4-q^{24}(q^4+1)(q^2+1)z^3+(q^{36}-q^{30}+q^{22}+q^{20}+2q^{18}+q^{16}+q^{14}-q^6+1)z^2-q^6(q^4+1)(q^2+1)z+q^6$,
$a^0_{12} = -q^3(q^{12}-1)(q^6+1)(1-z)(1+z)z$,
$a^0_{21} = -q^3\frac{q^6-1}{q^4-q^2+1}(1-z)(1+z)((q^{22}+q^{18})+(q^{40}-q^{38}+q^{36}-q^{34}-q^{30}-q^{26}-q^{20}-q^{14}-q^{10}-q^6+q^4-q^2+1)z+(q^{22}+q^{18})z^2)$,
$a^0_{22} = q^{30}-q^{24}(q^4+1)(q^2+1)z+(q^{36}-q^{30}+q^{22}+q^{20}+2q^{18}+q^{16}+q^{14}-q^6+1)z^2-q^6(q^4+1)(q^2+1)z^3+q^6z^4$.

4.4 Fusion construction

Let l be a positive integer and $\mathfrak{S}_l$ the symmetric group of order l. Let s_i be the simple reflection (the permutation of i and $i+1$). Let $l(w)$ be the length of $w \in \mathfrak{S}_l$. Then for any $w \in \mathfrak{S}_l$, we can define $R_w(x_1,\dots,x_l) : V_{x_1}\otimes\cdots\otimes V_{x_l} \to V_{x_{w(1)}}\otimes\cdots\otimes V_{x_{w(l)}}$ as follows:

$$R_1(x_1,\dots,x_l) = 1.$$

$$R_{s_i}(x_1,\dots,x_l) = \left(\otimes_{j<i}\mathrm{id}_{V_{x_j}}\right)\otimes R(x_i,x_{i+1})\otimes\left(\otimes_{j>i+1}\mathrm{id}_{V_{x_j}}\right).$$

For w, w' with $l(ww') = l(w)+l(w')$,

$$R_{ww'}(x_1,\dots,x_l) = R_{w'}(x_{w(1)},\dots,x_{w(l)})\circ R_w(x_1,\dots,x_l).$$

Fix $r \in \mathbf{Z}_{>0}$. For each $l \in \mathbf{Z}_{>0}$, we put

$$R_l = R_{w_0}\left(q^{r(l-1)},q^{r(l-3)},\dots,q^{-r(l-1)}\right) :$$

$$V_{q^{r(l-1)}}\otimes V_{q^{r(l-3)}}\otimes\cdots\otimes V_{q^{-r(l-1)}} \longrightarrow V_{q^{-r(l-1)}}\otimes V_{q^{-r(l-3)}}\otimes\cdots\otimes V_{q^{r(l-1)}},$$

where $w_0 \in \mathfrak{S}_l$ is the permutation given by $i \mapsto l+1-i$. Then R_l is a $U'_q(\mathfrak{g})$-linear homomorphism. We define

$$V_l = \mathrm{Im}R_l.$$

Hence taking $\Lambda_1 - 2\Lambda_0$ as λ_0 in Definition 3.16. By Proposition 4.3, we have

$$\varphi(z) = (1-q^{12}z)(1-q^{10}z)(1-q^8z)(1-q^6z).$$

Puting

$$r = 3.$$

We see that

$$\varphi(q^{2kr}) \text{ does not vanish for any } k > 0.$$

By Proposition 4.3, we see

$$\begin{aligned}
R(q^{2r})(u_{2\Lambda_1}) &= (1-q^{18})(1-q^{16})(1-q^{14})(1-q^{12}),\\
R(q^{2r})(u^1_{\Lambda_1}) &= \sum_{j=1}^{3} a^{\Lambda_1}_{1j} u^j_{\Lambda_1},\\
R(q^{2r})(u^1_0) &= \sum_{j=1}^{2} a^0_{1j} u^j_0,\\
R(q^{2r})(u_{2\Lambda_2}) &= 0,\\
R(q^{2r})(u^1_{\Lambda_1} - u^2_{\Lambda_1}) &= 0,\\
R(q^{2r})(u^3_{\Lambda_1}) &= 0,\\
R(q^{2r})(q^3(q^{12}-q^6+1)u^1_0 - (q^6+1)u^2_0) &= 0.
\end{aligned}$$

Then $N = \ker R(q^{2r})$ contains $u_{3\Lambda_2}$, $u_{2\Lambda_2}$, $u^1_{\Lambda_1} - u^2_{\Lambda_1}$, $u^3_{\Lambda_1}$ and $q^3(q^{12}-q^6+1)u^1_0 - (q^6+1)u^2_0$. Therefore by (4.2.1), we have

$$\begin{aligned}
N \;\cong\;& U_q(G_2)u_{3\Lambda_2} \oplus U_q(G_2)u_{2\Lambda_2} \oplus U_q(G_2)u^3_{\Lambda_1}\\
&\oplus U_q(G_2)(u^1_{\Lambda_1} - u^2_{\Lambda_1}) \oplus U_q(G_2)(q^3(q^{12}-q^6+1)u^1_0 - (q^6+1)u^2_0).
\end{aligned}$$

Then we have

$$\begin{aligned}
\dim(V^{\otimes 2}/N) &= \dim\left(\bigoplus_{j=0}^{2} V(j\Lambda_1)\right)\\
&= \sharp\left\{\boxed{b}\otimes\boxed{a} \;\middle|\; \boxed{a}\boxed{b} \in B(2\Lambda_1)\right\}\\
&\quad +\sharp\left\{\boxed{a}\otimes\boxed{\cdot} \;\middle|\; \boxed{a} \in B(\Lambda_1)\right\} + \sharp\left\{\boxed{\cdot}\otimes\boxed{\cdot}\right\}
\end{aligned}$$

where $a, b \in \{1, \ldots, 6, 0_1, 0_2, \overline{6}, \ldots, \overline{1}\}$.
Hence,

$$\begin{aligned}
&\dim\left(V^{\otimes l} \Big/ \sum V^i \otimes N \otimes V^{\otimes(l-2-i)}\right)\\
&\quad= \sharp\left\{\boxed{b_n}\otimes\cdots\otimes\boxed{b_1}\otimes\boxed{\cdot}\otimes\cdots\otimes\boxed{\cdot} \;\middle|\; \begin{array}{l}\boxed{b_i}\boxed{b_{i+1}} \in B(2\Lambda_1)\\ \boxed{\cdot}\boxed{b_1} \in B(\Lambda_1)\end{array},\ 0 \le i < n \le l\right\}\\
&\quad= \dim\left(\bigoplus_{j=0}^{l} V(j\Lambda_1)\right).
\end{aligned}$$

By (3.3.10) of [9], V_l is the quotient of $V^{\otimes l} \Big/ \sum_{i=0}^{l-2} V^{\otimes i} \otimes N \otimes V^{(l-2-i)}$.

Therefore we have,

$$\dim(V_l)_\lambda \le \sum_{j=0}^{l} \dim V(j(\Lambda_1 - 2\Lambda_0))_\lambda.$$

We set $i_0 = 0$, then $\langle h_1, \lambda_0\rangle = 1$, $\langle h_2, \lambda_0\rangle = 0$, $\{i_1\} = \{1\}$, $\langle h_0, \lambda_0\rangle = -2$. Therefore applying Proposition 3.21, we obtain the following results.

Proposition 4.4

1. V_l has a crystal pseudobase.
2. $V_l \cong \bigoplus_{j=0}^{l}(V(j(\Lambda_1 - 2\Lambda_0)))$ as a $U_q(G_2)$-module.

5 Perfectness of the crystals

The algebra $U_q(G_2^{(1)})$ is $\mathbf{Q}(q)$-algebra generated by $\{e_i, f_i, t_i, t_i^{-1}\ (i \in I = \{0,1,2\})\}$. Let $U_q((G_2^{(1)})_J)$ be a $\mathbf{Q}(q)$-algebra generated by $\{e_i, f_i, t_i, t_i^{-1}\ (i \in J \subset I)\}$.
The algebras $U_q((G_2^{(1)})_{\{1,2\}})$ and $U_q((G_2^{(1)})_{\{0,1\}})$ are isomorphic to $U_q(G_2)$ and $U_q(A_2)$ respectively. Let $J_0 = \{1,2\}$ and $J_2 = \{1,0\}$ be the index set of A_2 and G_2, respectively. We define $\imath_i : J_i \to I$ by $\imath_i(j) = j$ $(i = 0, 2)$. In order to show that the crystal B^l is perfect, we have to show the following conditions.

(1) There exists a crystal base of $U_q(G_2)$ isomorphic to $\imath_0^*(B^l)$.
There exists a crystal base of $U_q(A_2)$ isomorphic to $\imath_2^*(B^l)$. Crystal base B^l is connected.

(2) For any $b \in B^l$, $\langle c, \varphi(b)\rangle \geq l$.

(3) The maps ε and φ:$B_l = \{b \in B^l \mid \langle c, \varepsilon(b)\rangle = l\} \longrightarrow (P_{cl}^{+})_l = \{\lambda \in \sum \mathbf{Z}_{\geq 0}\Lambda_i \mid \langle c, \lambda\rangle = 1\}$ are bijective.

(4) Crystal graph of $B^l \otimes B^l$ is connected.

At first we construct crystal B^l which satisfies condition (1) (§5.2, §5.3). Then we prove conditions (2)(3) in §5.4, and (4) in §5.5.

5.1 Preparation

Let $B^{A_2}(\lambda)$ (resp. $B^{G_2}(\lambda)$) be a crystal base of $U_q(A_2)$ (resp. $U_q(G_2)$) with the highest weight λ. We write $\left\llbracket \llbracket b_1 \rrbracket \cdots \llbracket b_n \rrbracket \right\rrbracket$ instead of Young tableau $\begin{array}{|c|c|c|}\hline b_1 & \cdots & b_n \\ \hline\end{array}$. We introduce following notations:

$$\begin{aligned}
\llbracket a \rrbracket^k &= \overbrace{\llbracket a \rrbracket \cdots \llbracket a \rrbracket}^{k} \quad (a = 1, \ldots, 6, 0_1, 0_2, \overline{6}, \ldots, \overline{1}),\\
\llbracket \mathcal{C} \rrbracket^k &= \tilde{f}_1^k \llbracket 6 \rrbracket^k = \begin{cases} \llbracket 6 \rrbracket^{\frac{k}{2}} \llbracket \overline{6} \rrbracket^{\frac{k}{2}}, & k \equiv 0 \pmod 2,\\ \llbracket 6 \rrbracket^{[\frac{k}{2}]} \llbracket 0_2 \rrbracket \llbracket \overline{6} \rrbracket^{[\frac{k}{2}]}, & k \equiv 1 \pmod 2,\end{cases}\\
\llbracket \mathcal{W} \rrbracket^k &= \tilde{f}_2^k \llbracket 2 \rrbracket^k = \begin{cases} \llbracket 2 \rrbracket^{\frac{2k}{3}} \llbracket 6 \rrbracket^{\frac{k}{3}} & k \equiv 0 \pmod 3,\\ \llbracket 2 \rrbracket^{2[\frac{k}{3}]} \llbracket 3 \rrbracket \llbracket 6 \rrbracket^{[\frac{k}{3}]} & k \equiv 1 \pmod 3,\\ \llbracket 2 \rrbracket^{2[\frac{k}{3}]+1} \llbracket 4 \rrbracket \llbracket 6 \rrbracket^{[\frac{k}{3}]} & k \equiv 2 \pmod 3,\end{cases}\\
\llbracket \overline{\mathcal{W}} \rrbracket^k &= \tilde{e}_2^k \llbracket \overline{2} \rrbracket^k = \begin{cases} \llbracket \overline{6} \rrbracket^{\frac{k}{3}} \llbracket \overline{2} \rrbracket^{\frac{2k}{3}} & k \equiv 0 \pmod 3,\\ \llbracket \overline{6} \rrbracket^{[\frac{k}{3}]} \llbracket \overline{3} \rrbracket \llbracket \overline{2} \rrbracket^{2[\frac{k}{3}]} & k \equiv 1 \pmod 3,\\ \llbracket \overline{6} \rrbracket^{[\frac{k}{3}]} \llbracket \overline{4} \rrbracket \llbracket \overline{2} \rrbracket^{2[\frac{k}{3}]+1} & k \equiv 2 \pmod 3.\end{cases}
\end{aligned}$$

Then we easily have

$$\llbracket \mathcal{C} \rrbracket^k = \llbracket 6 \rrbracket \llbracket \mathcal{C} \rrbracket^{k-2} \llbracket \overline{6} \rrbracket, \tag{5.1.1}$$

$$\llbracket \mathcal{W} \rrbracket^k = \llbracket 2 \rrbracket^2 \llbracket \mathcal{W} \rrbracket^{k-3} \llbracket 6 \rrbracket, \tag{5.1.2}$$

$$\llbracket \overline{\mathcal{W}} \rrbracket^k = \llbracket \overline{6} \rrbracket \llbracket \overline{\mathcal{W}} \rrbracket^{k-3} \llbracket \overline{2} \rrbracket^2, \tag{5.1.3}$$

$$\tilde{f}_1\llbracket\mathcal{C}\rrbracket^{k} \;=\; \llbracket\mathcal{C}\rrbracket^{k-1}\llbracket\,\overline{6}\,\rrbracket\;, \tag{5.1.4}$$

$$\tilde{e}_2\llbracket\mathcal{W}\rrbracket^{k} \;=\; \llbracket\,2\,\rrbracket\,\llbracket\mathcal{W}\rrbracket^{k-1}\;, \tag{5.1.5}$$

$$\tilde{e}_2\llbracket\overline{\mathcal{W}}\rrbracket^{k} \;=\; \llbracket\,\overline{6}\,\rrbracket\,\llbracket\overline{\mathcal{W}}\rrbracket^{k-2}\llbracket\,2\,\rrbracket\;. \tag{5.1.6}$$

Proposition 5.1 *For* $\llbracket\mathcal{C}\rrbracket^{k}$, $\llbracket\mathcal{W}\rrbracket^{k}$, $\llbracket\overline{\mathcal{W}}\rrbracket^{k}$ $(k\in\mathbf{Z}_{\geq 0})$, *We have*

$$\mathrm{Red}_2\Big(\llbracket\mathcal{C}\rrbracket^{k}\Big) \;=\; u_0, \tag{5.1.7}$$

$$\mathrm{Red}_1\Big(\llbracket\mathcal{W}\rrbracket^{k}\Big) \;=\; u_0, \tag{5.1.8}$$

$$\mathrm{Red}_1\Big(\llbracket\overline{\mathcal{W}}\rrbracket^{k}\Big) \;=\; u_0. \tag{5.1.9}$$

Proof We prove (5.1.7). By Definition 3.4 and (3.1.1) we have

$$\Psi_2\Big(\llbracket\mathcal{C}\rrbracket^{2k}\Big) = \Psi_2\Big(\llbracket\,6\,\rrbracket^{k}\llbracket\,\overline{6}\,\rrbracket^{k}\Big) = u_+^{3k}u_-^{3k},$$

$$\mathrm{Red}_2\Big(\llbracket\mathcal{C}\rrbracket^{2k}\Big) = u_0,$$

$$\Psi_2\Big(\llbracket\mathcal{C}\rrbracket^{2k+1}\Big) = \Psi_2\Big(\llbracket\,6\,\rrbracket^{k}\llbracket 0_2\rrbracket\,\llbracket\,\overline{6}\,\rrbracket^{k}\Big) = u_+^{3k}u_0u_-^{3k},$$

$$\mathrm{Red}_2\Big(\llbracket\mathcal{C}\rrbracket^{2k+1}\Big) = u_0.$$

In a similar way, we can prove (5.1.8), (5.1.9). □

We use the following notations for $b\in B^{A_2}(m\Lambda_1+n\Lambda_2)$,

$$\begin{aligned} b \;&=\; \begin{array}{|c|c|c|c|c|c|}\hline b_1^{m+n} & \cdots & b_1^{m+1} & b_1^{m} & \cdots & b_1^{1} \\ \hline b_2^{m+n} & \cdots & b_2^{m+1} & \multicolumn{3}{c}{} \\ \cline{1-3}\end{array} \;\in B^{A_2}(m\Lambda_1+n\Lambda_2) \\ &=\; \Big\llbracket\,\llbracket b_1^{m+n},b_2^{m+n}\rrbracket\;\cdots\llbracket b_1^{m+1},b_2^{m+1}\rrbracket\,\llbracket b_1^{m}\rrbracket\;\cdots\llbracket b_1^{1}\rrbracket\,\Big\rrbracket. \end{aligned} \tag{5.1.10}$$

The highest weight element is given by

$$\begin{array}{|c|c|c|c|c|c|}\hline 1 & \cdots & 1 & 1 & \cdots & 1 \\ \hline 2 & \cdots & 2 & \multicolumn{3}{c}{} \\ \cline{1-3}\end{array} \;=\; \Big\llbracket\,\llbracket 1,2\rrbracket^{n}\,\llbracket\,1\,\rrbracket^{m}\,\Big\rrbracket.$$

Proposition 5.2 *Fix* $l\in\mathbf{Z}_{>0}$*. Take* i*,* j*,* p *such that* $0\leq i\leq\left[\frac{l}{2}\right]$*,* $i\leq j\leq l-i$*,* $0\leq p\leq j$*. Let* $\overline{b}\in B^{A_2}((i+j)\Lambda_1+l\Lambda_2)$ *be the highest weight element. Let* $\underline{b}\in B^{A_2}((i+j)\Lambda_1+l\Lambda_2)$ *be the lowest weight element. Then we have*

$$\varepsilon_2\left(\tilde{f}_1^{l+j}\tilde{f}_2^{l}\overline{b}\right) \;=\; 0, \tag{5.1.11}$$

$$\varphi_2\left(\tilde{f}_1^{l+j}\tilde{f}_2^{l}\overline{b}\right) \;=\; j, \tag{5.1.12}$$

$$\tilde{f}_2^{p}\tilde{f}_1^{l+j}\tilde{f}_2^{l}\overline{b} \;=\; \tilde{f}_1^{l+j-p}\tilde{f}_2^{l+p}\tilde{f}_1^{p}\overline{b}, \tag{5.1.13}$$

$$\tilde{f}_1^{l+j-p}\tilde{f}_2^{l+p}\tilde{f}_1^{p}\overline{b} \;=\; \tilde{e}_1^{i}\tilde{e}_2^{l-p}\underline{b}. \tag{5.1.14}$$

Proof By Proposition 3.13, we have

$$\overline{b} = \left\llbracket \llbracket 1,2 \rrbracket^{l} \llbracket 1 \rrbracket^{i+j} \right\rrbracket,$$

$$\underline{b} = \left\llbracket \llbracket 2,3 \rrbracket^{l} \llbracket 3 \rrbracket^{i+j} \right\rrbracket.$$

We prove (5.1.11), (5.1.12). Using Proposition 3.9, we have

$$\begin{aligned} \Psi_2(\overline{b}) &= u_0^{i+j}(u_0u_+)^l, \\ \mathrm{Red}_2\left(\overline{b}\right) &= u_+^l. \end{aligned}$$

Then we have

$$\tilde{f}_2^l\overline{b} = \left\llbracket \llbracket 1,3 \rrbracket^{l} \llbracket 1 \rrbracket^{i+j} \right\rrbracket.$$

Similarly, we have

$$\tilde{f}_1^{l+j}\tilde{f}_2^l\overline{b} = \left\llbracket \llbracket 1,3 \rrbracket^{i} \llbracket 2,3 \rrbracket^{l-i} \llbracket 2 \rrbracket^{i+j} \right\rrbracket.$$

Using Proposition 3.9, we have

$$\begin{aligned} \Psi_2\left(\left\llbracket \llbracket 1,3 \rrbracket^{i} \llbracket 2,3 \rrbracket^{l-i} \llbracket 2 \rrbracket^{i+j} \right\rrbracket\right) &= u_+^{i+j}(u_+u_-)^{l-i}(u_0u_-)^i, \\ \mathrm{Red}_2\left(\left\llbracket \llbracket 1,3 \rrbracket^{i} \llbracket 2,3 \rrbracket^{l-i} \llbracket 2 \rrbracket^{i+j} \right\rrbracket\right) &= u_+^j. \end{aligned}$$

Therefore, we have

$$\varepsilon_2\left(\left\llbracket \llbracket 1,3 \rrbracket^{i} \llbracket 2,3 \rrbracket^{l-i} \llbracket 2 \rrbracket^{i+j} \right\rrbracket\right) = 0,$$

$$\varphi_2\left(\left\llbracket \llbracket 1,3 \rrbracket^{i} \llbracket 2,3 \rrbracket^{l-i} \llbracket 2 \rrbracket^{i+j} \right\rrbracket\right) = j.$$

Relations (5.1.13), (5.1.14) can be proved by using

$$\tilde{f}_1^{l+j-p}\tilde{f}_2^{l+p}\tilde{f}_1^p\overline{b} = \left\llbracket \llbracket 1,3 \rrbracket^{i} \llbracket 2,3 \rrbracket^{l-i} \llbracket 2 \rrbracket^{i+j-p} \llbracket 3 \rrbracket^{p} \right\rrbracket.$$

□

Proposition 5.3 Let $\overline{b}$ be the highest weight element of the crystal base $B^{A_2}(k\Lambda_1+j\Lambda_2)$ of $U_q(A_2)$. Then for $0 \leq q < p \leq j$ we have,

$$\tilde{f}_1\tilde{f}_2\left(\tilde{f}_1^q\tilde{f}_2^p\overline{b}\right) = \tilde{f}_2\tilde{f}_1\left(\tilde{f}_1^q\tilde{f}_2^p\overline{b}\right).$$

Proof By (5.1.10), we have

$$\overline{b} = \left\llbracket \llbracket 1,2 \rrbracket^{j} \llbracket 1 \rrbracket^{k} \right\rrbracket.$$

Using Proposition 3.9, we have

$$\begin{aligned} \Psi_2(\overline{b}) &= u_0^k(u_0u_+)^j, \\ \mathrm{Red}_2\left(\overline{b}\right) &= u_+^j. \end{aligned}$$

Therefore we have

$$\tilde{f}_2^p\overline{b} = \left\llbracket \llbracket 1,2 \rrbracket^{j-p} \llbracket 1,3 \rrbracket^{p} \llbracket 1 \rrbracket^{k} \right\rrbracket.$$

Using Proposition 3.9 again, we have

$$\Psi_1\left(\tilde{f}_2^p\overline{b}\right) = u_+^k(u_+u_-)^p(u_+u_-)^{j-p}.$$

Thus we have

$$\tilde{f}_1^q \tilde{f}_2^p \overline{b} = \begin{cases} \left\llbracket \llbracket 1,2 \rrbracket^{j-p} \llbracket 1,3 \rrbracket^{p} \llbracket 1 \rrbracket^{k-q} \llbracket 2 \rrbracket^{q} \right\rrbracket & (k > q), \\ \left\llbracket \llbracket 1,2 \rrbracket^{j-p} \llbracket 1,3 \rrbracket^{p-q+k} \llbracket 2,3 \rrbracket^{q-k} \llbracket 2 \rrbracket^{k} \right\rrbracket & (k \leq q). \end{cases}$$

In a similar way we have,

$$\tilde{f}_2 \tilde{f}_1 \left(\tilde{f}_1^q \tilde{f}_2^p \overline{b} \right) = \tilde{f}_1 \tilde{f}_2 \left(\tilde{f}_1^q \tilde{f}_2^p \overline{b} \right) = \begin{cases} \left\llbracket \llbracket 1,2 \rrbracket^{j-p} \llbracket 1,3 \rrbracket^{p} \llbracket 1 \rrbracket^{k-q-1} \llbracket 2 \rrbracket^{q} \llbracket 3 \rrbracket \right\rrbracket & (k \geq q), \\ \left\llbracket \llbracket 1,2 \rrbracket^{j-p} \llbracket 1,3 \rrbracket^{p-q+k-1} \llbracket 2,3 \rrbracket^{q-k+1} \llbracket 2 \rrbracket^{k-1} \llbracket 3 \rrbracket \right\rrbracket & (k < q). \end{cases}$$

In a similar way, we have the following Proposition.

Proposition 5.4 *Let $\overline{b}$ be the highest weight element of $B^{A_2}(k\Lambda_1 + j\Lambda_2)$. For $0 \leq p \leq j$, $0 \leq s \leq k$, we have*

$$\tilde{e}_1 \tilde{f}_2^s \tilde{f}_1^{p+s+1} \tilde{f}_2^p \overline{b} = \tilde{f}_2^s \tilde{f}_1^{p+s} \tilde{f}_2^p \overline{b} \neq \tilde{f}_2^{s+1} \tilde{f}_1^{p+s} \tilde{f}_2^{p-1} \overline{b}, \tag{5.1.15}$$

$$\tilde{e}_1 \tilde{f}_2^{s+1} \tilde{f}_1^{p+s+1} \tilde{f}_2^p \overline{b} = \tilde{f}_2^{s+2} \tilde{f}_1^{p+s} \tilde{f}_2^{p-1} \overline{b} \neq \tilde{f}_2^{s+1} \tilde{f}_1^{p+s} \tilde{f}_2^p \overline{b}. \tag{5.1.16}$$

Proposition 5.5 *Let $\overline{b}$ be the highest weight element of crystal base $B^{A_2}(k\Lambda_1 + j\Lambda_2)$. For an element $\tilde{f}_1^q \tilde{f}_2^p \overline{b}$ $(p \leq q \leq p+k)$, we have*

$$\varphi_2 \left(\tilde{f}_1^q \tilde{f}_2^p b \right) = j + q - 2p. \tag{5.1.17}$$

Proof Using Proposition 3.9 we have,

$$\mathrm{Red}_2 \left(\tilde{f}_1^q \tilde{f}_2^p \overline{b} \right) = u_+^{j+p-2q}.$$

Proposition 5.6 For $b \in B^{G_2}(l\Lambda)$ such that $\tilde{f}_2 b \neq 0$, there exists unique $k' \in \{0, \ldots, \varphi_2(b) - 1\}$ such that

$$\varphi_1(\tilde{f}_2^{(k+1)}(b)) = \begin{cases} \varphi_1(\tilde{f}_2^{(k)}(b)) & k \leq k', \\ \varphi_1(\tilde{f}_2^{(k)}(b)) + 1 & k > k'. \end{cases}$$

where $k = 0, \ldots, \varphi_2(b) - 1$.

Proof Put $b = b_1 \otimes \cdots \otimes b_l$, and $r_1 r_2 \cdots r_{\varepsilon_1(b)+\varphi_1(b)} = \mathrm{Red}_{\mathfrak{n}}(b)$. Let k_1 be $r_{\varepsilon_1(b)}$ if $\varepsilon_1(b) \neq 0$, 0 if $\varepsilon_1(b) = 0$. By Proposition 3.3, there exists integers $k_2 \in \mathbf{Z}_{\geq 0}$ such that

$$\tilde{f}_2 b = b_1 \otimes \cdots \otimes \tilde{f}_2 b_{k_2} \otimes \cdots \otimes b_l.$$

First we consider $\Psi_1(\tilde{f}_2 b_{k_2})$, using u_+, u_- in Defnition 3.4.

$$\text{if } b_{k_2} = \llbracket 2 \rrbracket, \llbracket 5 \rrbracket, \llbracket \overline{6} \rrbracket, \llbracket \overline{4} \rrbracket, \ \text{ then } \Psi_1(b_{k_2}) = u_-^n,\ \Psi_1(\tilde{f}_2 b_{k_2}) = u_-^{n-1}\ (n = 1, 2). \tag{5.1.18}$$

$$\text{if } b_{k_2} = \llbracket 3 \rrbracket, \llbracket 4 \rrbracket, \llbracket 0_1 \rrbracket, \llbracket \overline{3} \rrbracket, \ \text{ then } \Psi_1(b_{k_2}) = u_+^{n'},\ \Psi_1(\tilde{f}_2 b_{k_2}) = u_+^{n'+1}\ (n = 0, 1). \tag{5.1.19}$$

We consider $\Psi_1(\tilde{f}_2 b)$. By Remark 3.10, we have $\mathrm{Red}_1(b) = u_-^{\varepsilon_1(b)} u_+^{\varphi_1(b)}$.

The case of $1 \leq k_2 \leq k_1$.
If (5.1.18), we see that

$$\mathrm{Red}_1(\tilde{f}_2 b) = u_-^{\varepsilon_1(b)-1} u_+^{\varphi_1(b)}. \tag{5.1.20}$$

If (5.1.19), we see that the increased u_+ is paired with a neighboring u_- and reduced. Then we have (5.1.20).

The case of $k_1+1 \leq k_2 \leq l$.
If (5.1.18), the decreased u_- is paired with u_+ in $\Psi_1(b)$. Then we see

$$\mathrm{Red}_1(\tilde{f}_2 b) = u_-^{\varepsilon_1(b)} u_+^{\varphi_1(b)+1}. \tag{5.1.21}$$

If (5.1.19), we see (5.1.21).

Therefore, we have $\varphi_1(\tilde{f}_2 b)$ equals $\varphi_1(b)$ or $\varphi_1(b)+1$.

Put $\tilde{f}_2 b = b' = b'_1 \otimes \cdots \otimes b'_l$. If $\tilde{f}_2 b' \neq 0$, we have

$$\tilde{f}_2 b' = b'_1 \otimes \cdots \otimes \tilde{f}_2 b_{k'_2} \otimes \cdots \otimes b'_l,$$

where $k'_2 \in \mathbf{Z}_{\geq 0}$. Here the action of $\tilde{f}_2$ is given by changing the leftmost u_+ to u_- in $\mathrm{Red}_2(b)$ and $\mathrm{Red}_2(b') = u_-^{\varepsilon_1(b')} u_+^{\varphi_1(b')}$. Then we see that $k'_2 \geq k_2$.

Thus we have the proposition.

5.2 Construction of crystal base B^l of $U_q(G_2^{(1)})$

5.2.1 Structure of B^l

As we noted earlier, $U_q((G_2^{(1)})_{\{1,2\}})$ is isomorphic to $U_q(G_2)$ and $U_q((G_2^{(1)})_{\{1,0\}})$ is isomorphic to $U_q(A_2)$. Since the structure of $U_q(A_2)$ is simpler than that of $U_q(G_2)$, we define the actions of $\tilde{f}_0$ on $U_q(G_2)$ by expoiting operators on $U_q(A_2)$. In view of isomorphism between $U_q((G_2^{(1)})_{\{1,0\}})$ and $U_q(A_2)$, we write $\{1,0\}$ as index set of roots of $U_q(A_2)$. The affine crystal B^l for $U_q(G_2^{(1)})$ is constructed on $\mathcal{G}^l$ with $\tilde{f}_0^{\mathcal{G}}$.

Definition 5.7 *Fix $l \in \mathbf{Z}_{>0}$. We define crystal base $\mathcal{G}^l$ of $U_q(G_2)$ and crystal base $\mathcal{A}$ of $U_q(A_2)$ as follows.*

$$\begin{aligned}
\mathcal{G}^l &= \bigoplus_{n=0}^{l} B^{G_2}(n\Lambda_1),\\
\mathcal{A}_i &= \bigoplus_{i \leq j_1, j_0 \leq l-i} B^{A_2}, (j_1\Lambda_1 + j_0\Lambda_0)\\
\mathcal{A} &= \bigoplus_{i=0}^{[\frac{l}{2}]} \mathcal{A}_i.
\end{aligned}$$

We denote $\tilde{f}_i$, $\tilde{e}_i$, φ_i, ε_i, wt, wt_i $(i=1,0)$ on $\mathcal{A}$ by $\tilde{f}_i^{\mathcal{A}}$, $\tilde{e}_i^{\mathcal{A}}$, $\varphi_i^{\mathcal{A}}$, $\varepsilon_i^{\mathcal{A}}$, $\mathrm{wt}^{\mathcal{A}}$, $\mathrm{wt}_i^{\mathcal{A}}$ respectively. In a similar way, we denote $\tilde{f}_i$, $\tilde{e}_i$, φ_i, ε_i, wt, wt_i $(i=1,2)$ on $\mathcal{G}$ by $\tilde{f}_i^{\mathcal{G}}$, $\tilde{e}_i^{\mathcal{G}}$, $\varphi_i^{\mathcal{G}}$, $\varepsilon_i^{\mathcal{G}}$, $\mathrm{wt}^{\mathcal{G}}$, $\mathrm{wt}_i^{\mathcal{G}}$ respectively.

Proposition 5.8

$$\sharp\mathcal{G}^l = \sharp\mathcal{A}$$

Proof The proposition amounts to show

$$\sum_{n=0}^{l} \sharp B^{G_2}(n\Lambda_1) = \sum_{i=0}^{[\frac{l}{2}]} \sum_{i \leq j_1, j_2 \leq l-i} \sharp B^{A_2}(j_1\Lambda_1 + j_2\Lambda_2). \tag{5.2.1}$$

Let Δ^+ be the set of positive roots, put $\delta = \frac{1}{2}\sum_{\beta\in\Delta^+}\beta$, and let $d(\Lambda)$ be the dimension of finite dimensinal irreducible representation with highest weight Λ. Then we have

$$d(\Lambda) = \prod_{\beta\in\Delta^+} \frac{(\Lambda+\delta,\beta)}{(\delta,\beta)}$$

by [13]. Therefore, we have

$$\sharp B^{G_2}(n\Lambda_1) = (n+1)\left(\frac{1}{2}n+1\right)\left(\frac{2}{3}n+1\right)\left(\frac{3}{4}n+1\right)\left(\frac{3}{5}n+1\right),$$

$$\sharp B^{A_2}(n_1\Lambda_1+n_2\Lambda_2)=\frac{1}{2}(n_1+1)(n_2+1)(n_1+n_2+2).$$

By calculation, we see that

$$\sharp B^{G_2}(l\Lambda_1)=\sum_{i=0}^{\left[\frac{l}{2}\right]}\left(\sum_{k=i}^{l-i-1}\sharp B^{A_2}(k\Lambda_1+(l-i)\Lambda_2)+\sum_{j=i}^{l-i}\sharp B^{A_2}((l-i)\Lambda_1+j\Lambda_0)\right). \tag{5.2.2}$$

Thus, we have (5.2.1). $\square$

By Proposition 5.8, we can construct one-to-one correspondence between $\mathcal{G}^l$ and $\mathcal{A}$ as sets.

Let $\overline{b}^{l,i}_{(j_1,j_0)}\in\mathcal{A}_i$ be the heighest weight element with weight $j_1\Lambda_1+j_0\Lambda_0$. Let $\underline{b}^{l,i}_{(-j_1,-j_0)}\in\mathcal{A}_i$ be the lowest weight element with weight $-j_1\Lambda_1-j_0\Lambda_0$. Elements $\overline{b}^{l,i}_{(j_1,j_0)}$ and $\underline{b}^{l,i}_{(-j_0,-j_1)}$ satisfy

$$\overline{b}^{l,i}_{(j_1,j_0)}=\left(\tilde{e}^{\mathcal{A}}_0\right)^{j_0}\left(\tilde{e}^{\mathcal{A}}_1\right)^{j_1+j_0}\left(\tilde{e}^{\mathcal{A}}_0\right)^{j_1}\underline{b}^{l,i}_{(-j_0,-j_1)}.$$

For a Lie algebra of type A_2, it is known that all elements b in the crystal base $B^{A_2}(j_i\Lambda_1+j_0\Lambda_0)$ are uniquely expressed as

$$b=\left(\tilde{f}^{\mathcal{A}}_0\right)^r\left(\tilde{f}^{\mathcal{A}}_1\right)^q\left(\tilde{f}^{\mathcal{A}}_0\right)^p\overline{b}^{l,i}_{(j_1,j_0)}\ (0\le p\le j_0, p\le q\le p+j_1, 0\le r\le j_0+q-2p), \tag{5.2.3}$$

It is possible that there exists crystal bases with same highest weight in $\mathcal{A}_i$ and $\mathcal{A}_{i'}$ $(i\neq i')$. To distinguish them, we define crystal base $B^i_{(j_1,j_0)}\subset\mathcal{A}_i$ by

$$B^i_{(j_1,j_0)}=\left\{\left(\tilde{f}^{\mathcal{A}}_0\right)^r\left(\tilde{f}^{\mathcal{A}}_1\right)^q\left(\tilde{f}^{\mathcal{A}}_0\right)^p\overline{b}^{l,i}_{(j_1,j_0)}\ \middle|\ 0\le p\le j_0,\ p\le q\le p+j_1,\ 0\le r\le j_0+q-2p\right\}. \tag{5.2.4}$$

By calculation, we have the following proposition.

Proposition 5.9 *For $B^i_{(k,j)}$ $(i<k, j\le l-i)$, we have*

$$\begin{aligned}&B^i_{(k,j)}\setminus\left(\left\{\left(\tilde{f}^{\mathcal{A}}_1\right)^q\left(\tilde{f}^{\mathcal{A}}_0\right)^p\overline{b}^{l,i}_{(k,j)}\ \middle|\ {}^{i<k,j\le l-i}_{0\le p\le j,0\le q\le p+k}\right\}\sqcup\left\{\left(\tilde{e}^{\mathcal{A}}_1\right)^q\left(\tilde{e}^{\mathcal{A}}_0\right)^p\underline{b}^{l,i}_{(-j,-k)}\ \middle|\ {}^{i<k,j\le l-i}_{0\le p\le j,0\le q\le p+k}\right\}\right)\\&\quad\cong B^{A_2}((k-1)\Lambda_1+(j-1)\Lambda_0).\end{aligned} \tag{5.2.5}$$

For $l\in\mathbf{Z}_{>0}$, we define

$$\begin{aligned}\mathcal{A}^{(l)}_i&=\left(\bigsqcup_{i\le k\le l-i}B^i_{(k,i)}\right)\bigsqcup\left(\bigsqcup_{i<j\le l-q}B^i_{(i,j)}\right)\\&\quad\bigsqcup\left(\bigsqcup_{\substack{i<k,j\le l-i\\0\le p\le j,0\le q\le p+k}}\left(\tilde{f}^{\mathcal{A}}_1\right)^q\left(\tilde{f}^{\mathcal{A}}_0\right)^p\overline{b}^{l,i}_{(k,j)}\right)\bigsqcup\left(\bigsqcup_{\substack{i<k,j\le l-i\\0\le p<k,0\le q\le p+j}}\left(\tilde{e}^{\mathcal{A}}_1\right)^q\left(\tilde{e}^{\mathcal{A}}_0\right)^p\underline{b}^{l,i}_{(-j,-k)}\right),\\\mathcal{A}^{(l)}&=\bigsqcup_{0\le i\le\left[\frac{l}{2}\right]}\mathcal{A}^{(l)}_i.\end{aligned}$$

Proposition 5.10 *For $l\in\mathbf{Z}_{>0}$, we have*

$$\sharp B^{G_2}(l\Lambda_1)=\sharp\mathcal{A}^{(l)}.$$

Proof By Proposition 5.9, we have

$$\begin{aligned}
\sharp B^{A_2}\left((l-i)\Lambda_1+(l-k)\Lambda_0\right) &= \sharp B^{i}_{(k,i)} \\
&\quad + \sum_{m=1}^{l-i-k} \sharp\left\{ \left(\tilde{f}_1^{\mathcal{A}}\right)^{q} \left(\tilde{f}_0^{\mathcal{A}}\right)^{p} \overline{b}^{\,l,i}_{(k+m,i+m)} \,\middle|\, \begin{array}{c} 0\le p\le i+m \\ 0\le q\le p+k+m \end{array}\right\} \\
&\quad + \sum_{m=1}^{l-i-k} \sharp\left\{ \left(\tilde{e}_1^{\mathcal{A}}\right)^{q} \left(\tilde{e}_0^{\mathcal{A}}\right)^{p} \underline{b}^{\,l,i}_{(-i-m,-k-m)} \,\middle|\, \begin{array}{c} 0\le p< k+m \\ 0\le q\le p+i+m \end{array}\right\}, \\
\sharp B^{A_2}\left((l-j)\Lambda_1+(l-i)\Lambda_0\right) &= \sharp B^{i}_{(i,j)} \\
&\quad + \sum_{m=1}^{l-i-j} \sharp\left\{ \left(\tilde{f}_1^{\mathcal{A}}\right)^{q} \left(\tilde{f}_0^{\mathcal{A}}\right)^{p} \overline{b}^{\,l,i}_{(i+m,j+m)} \,\middle|\, \begin{array}{c} 0\le p\le j+m \\ 0\le q\le p+i+m \end{array}\right\} \\
&\quad + \sum_{m=1}^{l-i-j} \sharp\left\{ \left(\tilde{e}_1^{\mathcal{A}}\right)^{q} \left(\tilde{e}_0^{\mathcal{A}}\right)^{p} \underline{b}^{\,l,i}_{(-j-m,-i-m)} \,\middle|\, \begin{array}{c} 0\le p< i+m \\ 0\le q\le p+j+m \end{array}\right\}.
\end{aligned}$$

By (5.2.2), we have

$$\sum_{i=0}^{\left[\frac{l}{2}\right]}\left(\sum_{k=i}^{l-i}\sharp B^{A_2}((l-k)\Lambda_1+(l-i)\Lambda_0)+\sum_{j=i+1}^{l-i}\sharp B^{A_2}((l-i)\Lambda_1+(l-j)\Lambda_0)\right)=\sharp B^{G_2}(l\Lambda_1).$$

□

By Proposition 5.9 and Proposition 5.10, we have the following lemma.

Lemma 5.11 *For* $l \in \mathbf{Z}_{>0}$ *we have*

$$\mathcal{G}^{l} \setminus \mathcal{A}^{(l)} \cong \mathcal{G}^{l-1}.$$

as crystal base of $U_q(A_2)$.

5.2.2 Operators $E_{\mathcal{A}}$ and $F_{\mathcal{A}}$ on $\mathcal{A}$

We are going to define operators $E_{\mathcal{A}}$ and $F_{\mathcal{A}}$ on $\mathcal{A}$ which satisfy properties

(C1) for $b,\ b' \in \mathcal{A}$, $E_{\mathcal{A}}b = b'$, if and only if $F_{\mathcal{A}}b' = b$,

(C2) for $b \in \mathcal{A}$, $E_{\mathcal{A}}\tilde{f}_0^{\mathcal{A}}(b) = \tilde{f}_0^{\mathcal{A}}E_{\mathcal{A}}(b)$,

(C3) for $b \in \mathcal{A}$, $\max\{m \mid F_{\mathcal{A}}^{m}b \neq 0\} - \max\{m' \mid E_{\mathcal{A}}^{m'}b \neq 0\} = -2\mathrm{wt}_1^{\mathcal{A}}(b) - \mathrm{wt}_0^{\mathcal{A}}(b)$.

Operators $E_{\mathcal{A}}$ and $F_{\mathcal{A}}$ are counterpart of $\tilde{e}_2^{\mathcal{G}}$ and $\tilde{f}_2^{\mathcal{G}}$, respectively. In view of Lemma 5.2.5 and (C2) we define $E_{\mathcal{A}}$ and $F_{\mathcal{A}}$ inductively. If $l = 0$, $\mathcal{A}$ is trivial. If $l > 0$, we use following relation for induction

$$E_{\mathcal{A}}\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q}\left(\tilde{f}_0^{\mathcal{A}}\right)^{p}\overline{b}^{\,l,i}_{(k,j)}\right) = \tilde{e}_0^{\mathcal{A}}\left(E_{\mathcal{A}}\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q-1}\left(\tilde{f}_0^{\mathcal{A}}\right)^{p}\overline{b}^{\,l-1,i}_{(k-1,j-1)}\right)\right). \tag{5.2.6}$$

Put $\mathcal{A}_{+}^{(l)} = \{b \mid \tilde{e}_0(b) = 0\}$. Using Proposition 3.9, we see that $\mathcal{A}_{+}^{(l)}$ is given by

$$\mathcal{A}_{+}^{(l)} = \left\{\left(\tilde{f}_1^{\mathcal{A}}\right)^{q}\left(\tilde{f}_0^{\mathcal{A}}\right)^{p}\overline{b}^{\,l,i}_{(k,j)} \,\middle|\, 0 \le i \le \left[\frac{l}{2}\right],\ i \le k, j \le l-i,\ 0 \le p \le j,\ p \le q \le p+k\right\}.$$

By (C1) and (C2), it is sufficient to define the operator $E_{\mathcal{A}}$ for $b \in \mathcal{A}_{+}^{(l)}$.

Definition 5.12 *We define $E_{\mathcal{A}}$ by*

$$E_{\mathcal{A}}\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^q\left(\tilde{f}_0^{\mathcal{A}}\right)^p\overline{b}^{l,i}_{(k,j)}\right)=\begin{cases}\left(\tilde{f}_1^{\mathcal{A}}\right)^q\left(\tilde{f}_0^{\mathcal{A}}\right)^p\overline{b}^{l,i}_{(k-1,j)}, & \text{(a1)}\\ \left(\tilde{f}_1^{\mathcal{A}}\right)^{q+1}\left(\tilde{f}_0^{\mathcal{A}}\right)^{p+1}\overline{b}^{l,i}_{(k,j+1)}, & \text{(a2)}\\ \left(\tilde{f}_1^{\mathcal{A}}\right)^{q+1}\left(\tilde{f}_0^{\mathcal{A}}\right)^{p}\overline{b}^{l,i+1}_{(k+1,j-1)}, & \text{(a3)}\\ \left(\tilde{f}_1^{\mathcal{A}}\right)^{q+1}\left(\tilde{f}_0^{\mathcal{A}}\right)^{p}\overline{b}^{l,i}_{(k+1,j-1)}, & \text{(a4)}\\ \left(\tilde{f}_1^{\mathcal{A}}\right)^{q+1}\left(\tilde{f}_0^{\mathcal{A}}\right)^{p}\overline{b}^{l,i-1}_{(k+1,j-1)}, & \text{(a5)}\\ 0, & \text{(a6)}\\ \tilde{e}_0^{\mathcal{A}}\left(E_{\mathcal{A}}\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q-1}\left(\tilde{f}_0^{\mathcal{A}}\right)^{p}\overline{b}^{l-1,i}_{(k-1,j-1)}\right)\right). & \text{(5.2.6)}\end{cases}$$

Conditions for (a1)-(a6) *and* (5.2.6) *are as follows:*

(1) *If $i<k\le l-i$, $i<j\le l-i$,*

  (i) *if $0\le q\le j-1-\left[\frac{j-k}{3}\right]$, $p=min(q,j)$, then the action of $E_{\mathcal{A}}$ is given by* (a1),

  (ii) *if $j<l-i$, $j-\left[\frac{j-k}{3}\right]\le q\le j+k$, $p=min(q,j)$, then the action of $E_{\mathcal{A}}$ is given by* (a2),

  (iii) *if $j=l-i$, $j-\left[\frac{j-k}{3}\right]\le q\le j+k$, $p=min(q,j)$, then the action of $E_{\mathcal{A}}$ is given by* (a6),

  (iv) *if $0\le p\le j-1$, $p+1\le q\le p+k$, then we use induction* (5.2.6).

(2) *If $k=i$, $i+2\le j\le l-i$,*

  (i) *if $0\le p\le j-1-\left[\frac{j-i}{3}\right]$, $p\le q\le p+i$ then the action of $E_{\mathcal{A}}$ is given by* (a3),

  (ii) *if $j<l-i$,$j-\left[\frac{j-i}{3}\right]\le p\le j$, $p\le q\le p+i$ then the action of $E_{\mathcal{A}}$ is given by* (a2),

  (iii) *if $j=l-i$, $j-\left[\frac{j-i}{3}\right]\le p\le j$, $p\le q\le p+i$ then the action of $E_{\mathcal{A}}$ is given by* (a6).

(3) *If $k=i$, $j=i+1$,*

  (i) *if $0\le p\le i$, $p\le q\le p+i$ then the action of $E_{\mathcal{A}}$ is given by* (a4),

  (ii) *if $p=i+1$, $p\le q\le p+i$ then the action of $E_{\mathcal{A}}$ is given by* (a2).

(4) *If $k=i$, $j=i$,*

  (i) *if $0\le p\le i-1$, $p\le q\le p+i$ then the action of $E_{\mathcal{A}}$ is given by* (a5),

  (ii) *if $p=i$, $i\le q\le 2i$ then the action of $E_{\mathcal{A}}$ is given by* (a2).

(5) *If $i+1\le k\le l-i$, $j=i$,*

  (i) *if $0\le p\le i$, $p\le q\le p-1-\left[\frac{i-k}{3}\right]$ then the action of $E_{\mathcal{A}}$ is given by* (a1),

  (ii) *if $p=i$, $p-\left[\frac{i-k}{3}\right]\le q\le p+k$ then the action of $E_{\mathcal{A}}$ is given by* (a2),

  (iii) *if $0\le p\le i-1$, $p-\left[\frac{i-k}{3}\right]\le q\le p+k$ then the action of $E_{\mathcal{A}}$ is given by* (a5).

By (C1) and definition of $E_{\mathcal{A}}$, we see the action of $F_{\mathcal{A}}$ on $\mathcal{A}_{+}^{(l)}$ is given by

$$F_{\mathcal{A}}\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^q\left(\tilde{f}_0^{\mathcal{A}}\right)^p\overline{b}_{(k,j)}^{l,i}\right)=\begin{cases}\tilde{f}_1^q\tilde{f}_0^p\overline{b}_{(k+1,j)}^{l,i}, & \text{(a1)}'\\ \tilde{f}_1^{q-1}\tilde{f}_0^{p-1}\overline{b}_{(k,j-1)}^{l,i}, & \text{(a2)}'\\ \tilde{f}_1^{q-1}\tilde{f}_0^p\overline{b}_{(k-1,j+1)}^{l,i-1}, & \text{(a3)}'\\ \tilde{f}_1^{q-1}\tilde{f}_0^p\overline{b}_{(k-1,j+1)}^{l,i}, & \text{(a4)}'\\ \tilde{f}_1^{q-1}\tilde{f}_0^p\overline{b}_{(k-1,j+1)}^{l,i+1}, & \text{(a5)}'\\ 0, & \text{(a6)}'\\ \tilde{e}_0^{\mathcal{A}}\left(F_{\mathcal{A}}\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q-1}\left(\tilde{f}_0^{\mathcal{A}}\right)^p\overline{b}_{(k-1,j-1)}^{l-1,i}\right)\right). & (5.2.6)'\end{cases}$$

As in the case of $E_{\mathcal{A}}$, we use induction (5.2.6)$'$. Conditions for (a1)$'$-(a6)$'$ are as follows:

(1) If $i < k \leq l-i$, $i < j \leq l-i$,

(i) if $k < l-i$, $0 \leq q \leq j-1-\left[\frac{j-k-1}{3}\right]$, $p=\min(q,j)$, then the action of $F_{\mathcal{A}}$ is given by (a1)$'$,

(ii) if $k = l-i$, $0 \leq q \leq j-1-\left[\frac{j-k-1}{3}\right]$, $p=\min(q,j)$, then the action of $F_{\mathcal{A}}$ is given by (a6)$'$,

(iii) if $j-\left[\frac{j-k-1}{3}\right] \leq q \leq j+k$, $p=\min(q,j)$, then the action of $F_{\mathcal{A}}$ is given by (a2)$'$,

(iv) if $0 \leq p \leq j-1$, $p+1 \leq q \leq p+k$, then we use induction (5.2.6)$'$.

(2) If $k=i$, $i+1 \leq j \leq l-i$,

(i) if $0 \leq p \leq j-1-\left[\frac{j-k-1}{3}\right]$, $q=p$ then the action of $F_{\mathcal{A}}$ is given by (a1)$'$,

(ii) if $0 \leq p \leq j-1-\left[\frac{j-k-1}{3}\right]$, $p+1 \leq q \leq p+k+1$ then the action of $F_{\mathcal{A}}$ is given by (a3)$'$,

(iii) if $j-\left[\frac{j-k-1}{3}\right] \leq p \leq j$, $p \leq q \leq p+k$, then the action of $F_{\mathcal{A}}$ is given by (a2)$'$.

(3) If $k=i$, $j=i$,

(i) if $0 \leq p \leq i$, $p=q$ then the action of $F_{\mathcal{A}}$ is given by (a2)$'$,

(ii) if $0 \leq p \leq i$, $p+1 \leq q \leq p+i$ then the action of $F_{\mathcal{A}}$ is given by (a3)$'$.

(4) If $k=i+1$, $j=i$,

(i) if $0 \leq p \leq i$, $p=q$ then the action of $F_{\mathcal{A}}$ is given by (a2)$'$,

(ii) if $0 \leq p \leq i$, $p+1 \leq q \leq p+i+1$ then the action of $F_{\mathcal{A}}$ is given by (a4)$'$.

(5) If $i+2 \leq k \leq l-i$, $j=i$,

(i) if $0 \leq p \leq i$, $p \leq q \leq p-1-\left[\frac{i-k-1}{3}\right]$ then the action of $F_{\mathcal{A}}$ is given by (a1)$'$,

(ii) if $0 \leq p \leq i$, $p-\left[\frac{i-k-1}{3}\right] \leq q \leq p+k$ then the action of $F_{\mathcal{A}}$ is given by (a5)$'$.

For $b \in \mathcal{A}$, we put

$$\varepsilon_{\mathcal{A}}(b)=\max\{m \mid F_{\mathcal{A}}^m b \neq 0\},$$

$$\varphi_{\mathcal{A}}(b)=\max\{m' \mid E_{\mathcal{A}}^{m'} b \neq 0\}.$$

We verify that $E_{\mathcal{A}}$ is well-defined. By the definition of $F_{\mathcal{A}}$, we see that for $b, b' \in \mathcal{A}_{+}^{(l)}$,

$$\text{if } E_{\mathcal{A}}(b)=E_{\mathcal{A}}(b') \neq 0, \text{ then } b=b'. \tag{5.2.7}$$

In order to verify (C2), we prove that $E_{\mathcal{A}}(\tilde{f}_0^{\mathcal{A}}(b)) \neq 0$ if and only if $\tilde{f}_0^{\mathcal{A}}(E_{\mathcal{A}}(b)) \neq 0$. For this, it is sufficient to prove that for $b \in A_+^{(l)}$,

$$\varphi_0(b) = \varphi_0(E_{\mathcal{A}} b). \tag{5.2.8}$$

Using Proposition 5.5, we have following formula in the case of (a1) of Definition 5.12,

$$\varphi_0\left(\tilde{f}_1^q \tilde{f}_0^p \overline{b}_{(k,j)}^{l,i}\right) = j + q - 2p,$$

$$\varphi_0\left(E_{\mathcal{A}} \tilde{f}_1^q \tilde{f}_0^p \overline{b}_{(k,j)}^{l,i}\right) = \varphi_0\left(\tilde{f}_1^q \tilde{f}_0^p \overline{b}_{(k-1,j)}^{l,i}\right) = j + q - 2p.$$

In a similar way we can prove other cases. Therefore we see that for $b \in \mathcal{A}$ such that $E_{\mathcal{A}}(b) \neq 0$, $\tilde{f}_0^{\mathcal{A}} b \neq 0$ we have $E_{\mathcal{A}}\left(\tilde{f}_0^{\mathcal{A}} b\right) \neq 0$. Similarly we can show that for $b \in \mathcal{A}$ such that $E_{\mathcal{A}}(b) = 0$ or $\tilde{f}_0^{\mathcal{A}} b = 0$ we have $E_{\mathcal{A}}\left(\tilde{f}_0^{\mathcal{A}} b\right) = 0$.

We verify (C3). By calcualtion, we see

$$\mathrm{wt}^{\mathcal{A}}\left(\tilde{f}_1^q \tilde{f}_0^p \overline{b}_{(k,j)}^{l,i}\right) = (k + p - 2q)\Lambda_1 + (j - 2p + q)\Lambda_0. \tag{5.2.9}$$

If the action of $E_{\mathcal{A}}$ is given by (a1), we have

$$\mathrm{wt}^{\mathcal{A}}\left(E_{\mathcal{A}} \tilde{f}_1^q \tilde{f}_0^p \overline{b}_{(k,j)}^{l,i}\right) = \mathrm{wt}^{\mathcal{A}}\left(\tilde{f}_1^q \tilde{f}_0^p \overline{b}_{(k-1,j)}^{l,i}\right) = (k - 1 + p - 2q)\Lambda_1 + (j - 2p + q)\Lambda_0.$$

Therefore, we have

$$-2\mathrm{wt}_1^{\mathcal{A}}(E_{\mathcal{A}} b) - \mathrm{wt}_0^{\mathcal{A}}(E_{\mathcal{A}} b) = -2\mathrm{wt}_1^{\mathcal{A}}(b) - \mathrm{wt}_0^{\mathcal{A}}(b) + 2. \tag{5.2.10}$$

Similarly we can show (5.2.10) when the action of $E_{\mathcal{A}}$ is given by (a2) - (a5).

We prove (C3) for $b = \left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}_{(l-i,j)}^{l,i} \in \mathcal{A}_+^{(l)}$ where $i \leq j \leq l - i$, $0 \leq p \leq j$, $p \leq q \leq p + l - i$. By Definition 5.12, b satisfies $F_{\mathcal{A}}(b) = 0$ and

$$\varepsilon_{\mathcal{A}}(b) = 2l - 2i + j - 3q. \tag{5.2.11}$$

1. If $0 \leq p = q \leq \left[\frac{i+j}{2}\right]$,
   The action of $E_{\mathcal{A}}$ on $\tilde{f}_1^p \tilde{f}_0^p \overline{b}_{(k,j)}^{l,i}$ $(i < k \leq l - i)$ is given by (a1). Then we have

   $$E_{\mathcal{A}}^{l-2i}(b) = \tilde{f}_1^p \tilde{f}_0^p \overline{b}_{(i,j)}^{l,i}.$$

   If $\left[\frac{j-i}{2}\right] > 0$, the action of $E_{\mathcal{A}}$ on $\tilde{f}_1^p \tilde{f}_0^p \overline{b}_{(i,j)}^{l,i}$ is given by (a3). We put $x = \frac{i+j}{2}$, $\hat{x} = \frac{i+j+1}{2}$. Then we have

   $$E_{\mathcal{A}}^{l-3i+[x]}(b) = \begin{cases} \tilde{f}_1^{p+\hat{x}-i-1} \tilde{f}_0^p \overline{b}_{(\hat{x}-1,\hat{x})}^{l,\hat{x}-1} & j - i \text{ is odd,} \\ \tilde{f}_1^{p+x-i} \tilde{f}_0^p \overline{b}_{(x,x)}^{l,x} & j - i \text{ is even.} \end{cases}$$

   If $i - j$ is odd, the action of $E_{\mathcal{A}}$ on $\tilde{f}_1^{p+\hat{x}-i-1} \tilde{f}_0^p \overline{b}_{((\hat{x}-1),\hat{x})}^{l,\hat{x}-1}$ is given by (a4). Then we have

   $$E_{\mathcal{A}}^{l-3i+[x]}(b) = \tilde{f}_1^{p+\hat{x}-i} \tilde{f}_0^p \overline{b}_{(\hat{x},(\hat{x}-1))}^{l,\hat{x}-1}.$$

   If $j - p - [\hat{x} - i] > 0$, then the action of $E_{\mathcal{A}}$ on $\tilde{f}_1^{p+x-i} \tilde{f}_0^p \overline{b}_{(x,x)}^{l,x}$, $\tilde{f}_1^{p+\hat{x}-i} \tilde{f}_0^p \overline{b}_{(\hat{x},\hat{x}-1)}^{l,\hat{x}-1}$ are given by (a5). Then we have

   $$E_{\mathcal{A}}^{l-2i+j-p}(b) = \tilde{f}_1^j \tilde{f}_0^p \overline{b}_{(j+i-p,p)}^{l,p}.$$

   Moreover, the action of $E_{\mathcal{A}}$ on $\tilde{f}_1^j \tilde{f}_0^p \overline{b}_{(j+i-p,p)}^{l,p}$ is given by (a2). Then we have

   $$E_{\mathcal{A}}^{2l-2i+j-3p}(b) = \tilde{f}_1^{l+j-2p} \tilde{f}_0^{l-p} \overline{b}_{(j+i-p,l-p)}^{l,p}. \tag{5.2.12}$$

   Finally, the action of $E_{\mathcal{A}}$ on $\tilde{f}_1^{l+j-2p} \tilde{f}_0^{l-p} \overline{b}_{(j+i-p,l-p)}^{l,p}$ is given by (a6). Then we have

   $$E_{\mathcal{A}}^{2l-2i+j-3p+1}(b) = 0.$$

2. If $\left[\frac{i+j}{2}\right] < p = q < \frac{2j+i}{3}$, we can calculate in a similar way:

$$\begin{array}{rcll} E_{\mathcal{A}}^{l-2i}(b) & = & \tilde{f}_1^p \tilde{f}_0^p \overline{b}_{(i,j)}^{l,i}, & \text{(a1)} \\ E_{\mathcal{A}}^{l-i+2j-3p}(b) & = & \tilde{f}_1^{2j+i-2p} \tilde{f}_0^p \overline{b}_{(2j+2i-3p,-j-i+3p)}^{l,2j+2i-3p}, & \text{(a3)} \\ E_{\mathcal{A}}^{2l-2i+j-3p}(b) & = & \tilde{f}_1^{l+j-2p} \tilde{f}_0^{l-i-j+p} \overline{b}_{(2j+2i-3p,l-2j-2i+3p)}^{l,2j+2i-3p}, & \text{(a2)} \\ E_{\mathcal{A}}^{2l-2i+j-3p+1}(b) & = & 0. & \text{(a6)} \end{array} \tag{5.2.13}$$

3. If $\frac{2j+i}{3} < p = q \leq j$ or $j = p < q$, we have

$$\begin{array}{rcll} E_{\mathcal{A}}^{l-i+2j-3q}(b) & = & \tilde{f}_1^q \tilde{f}_0^p \overline{b}_{(3q-2j,j)}^{l,i}, & \text{(a1)} \\ E_{\mathcal{A}}^{2l-2i+j-3q}(b) & = & \tilde{f}_1^{l-i-j+q} \tilde{f}_0^{l-i-j+p} \overline{b}_{(3q-2j,l-i)}^{l,i}, & \text{(a2)} \\ E_{\mathcal{A}}^{2l-2i+j-3q+1}(b) & = & 0. & \text{(a6)} \end{array} \tag{5.2.14}$$

4. If $p < j = i$, $p < q < y + j$, we have

$$\begin{array}{rcll} E_{\mathcal{A}}^{l-2i-3(p-q)}(b) & = & \tilde{f}_1^q \tilde{f}_0^p \overline{b}_{(i+3(q-p),i)}^{l,i}, & \text{(a1)} \\ E_{\mathcal{A}}^{l-i-3q+2p}(b) & = & \tilde{f}_1^{q+i-p} \tilde{f}_0^p \overline{b}_{(2i+3q-4p,l-p)}^{l,p}, & \text{(a5)} \\ E_{\mathcal{A}}^{2l-i-3q}(b) & = & \tilde{f}_1^{l+i-3p+q} \tilde{f}_0^{l-p} \overline{b}_{(2i+3q-4p,l-p)}^{l,p}, & \text{(a2)} \\ E_{\mathcal{A}}^{2l-i-3q+1}(b) & = & 0. & \text{(a6)} \end{array} \tag{5.2.15}$$

5. Other cases are reduced to $\mathcal{G}^{l-1}$, using (5.2.6).

Then for $b = \left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}_{(l-i,j)}^{l,i} \in \mathcal{A}_+^{(l)}$ where $i \leq j \leq l-i$, $0 \leq p \leq j$, $p \leq q \leq p+l-i$, we have

$$-2\mathrm{wt}_1^{\mathcal{A}}(b) - \mathrm{wt}_0^{\mathcal{A}}(b) = -(2l - 2i + j - 3q).$$

By (5.2.11), we see

$$\varphi_{\mathcal{A}}(b) - \varepsilon_{\mathcal{A}}(b) = -(2l - 2i + j - 3q).$$

Therefore by (5.2.10), we verify inductively that $E_{\mathcal{A}}$ satisfies (C3).

5.2.3 An involution on $\mathcal{A}$

Let $\underline{b}_{(j_1,j_0)}^{l,i}$ be the lowest weight element in $\mathcal{A}_i$ with weight $j_1\Lambda_1 + j_0\Lambda_0$. We define a map $\mathrm{C}_{\mathcal{A}} : \mathcal{A} \to \mathcal{A}$ by:

$$\begin{array}{rcl} \mathrm{C}_{\mathcal{A}}\left(\left(\tilde{f}_0^{\mathcal{A}}\right)^r \left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}_{(k,j)}^{l,i}\right) & = & \left(\tilde{e}_0^{\mathcal{A}}\right)^r \left(\tilde{e}_1^{\mathcal{A}}\right)^q \left(\tilde{e}_0^{\mathcal{A}}\right)^p \underline{b}_{(-k,-j)}^{l,i} \\ & = & \left(\tilde{f}_0^{\mathcal{A}}\right)^{j+q-2p-r} \left(\tilde{f}_1^{\mathcal{A}}\right)^{k+j-q} \left(\tilde{f}_0^{\mathcal{A}}\right)^{k-q+p} \overline{b}_{(j,k)}^{l,i}. \end{array} \tag{5.2.16}$$

where $0 \leq p \leq j$, $p \leq q \leq p+k$, $0 \leq r \leq j+q-2p$. It is easy to see that $\mathrm{C}_{\mathcal{A}}$ is an involution,

$$\mathrm{C}_{\mathcal{A}}\left(\mathrm{C}_{\mathcal{A}}(b)\right) = b \ \ (b \in \mathcal{A}).$$

Proposition 5.13 *For $b \in \mathcal{A}$, we have*

$$\mathrm{C}_{\mathcal{A}}\left(E_{\mathcal{A}}b\right) = F_{\mathcal{A}}(\mathrm{C}_{\mathcal{A}}(b)). \tag{5.2.17}$$

Proof It is sufficient to prove (5.2.17) for $b \in \mathcal{A}$ such that $\varepsilon_0^{\mathcal{A}}(b) = 0$.
We assume that for $b = \left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{\,l,i}_{(k,j)} \in \mathcal{A}$, k, j, p, q satisfy $i+1 \le k \le l-i$, $i+1 \le j \le l-i$, $0 \le q \le j-1-\left[\frac{j-k}{3}\right]$, $p = \min(q,j)$. Since this is the condition (1)(i) in Definition 5.12, the action of $E_{\mathcal{A}}$ is given by (a1). Then we have

$$\mathrm{C}_{\mathcal{A}}\left(E_{\mathcal{A}} b\right) = \left(\tilde{f}_0^{\mathcal{A}}\right)^{j+q-2p} \left(\tilde{f}_1^{\mathcal{A}}\right)^{k+j-q-1} \left(\tilde{f}_0^{\mathcal{A}}\right)^{k-q+p-1} \overline{b}^{\,l,i}_{(j,k-1)}.$$

We consider the action of $F_{\mathcal{A}}$ on $\mathrm{C}_{\mathcal{A}}(b)$. By applying the involution, we have

$$\mathrm{C}_{\mathcal{A}}(b) = \left(\tilde{f}_0^{\mathcal{A}}\right)^{j+q-2p} \left(\tilde{f}_1^{\mathcal{A}}\right)^{k+j-q} \left(\tilde{f}_0^{\mathcal{A}}\right)^{k-q+p} \overline{b}^{\,l,i}_{(j,k)}.$$

Thus the action of $F_{\mathcal{A}}$ on $\mathrm{C}_{\mathcal{A}}(b)$ is given by (a2)′. Therefore we have

$$F_{\mathcal{A}}\left(\mathrm{C}_{\mathcal{A}}(b)\right) = \left(\tilde{f}_0^{\mathcal{A}}\right)^{j+q-2p} \left(\tilde{f}_1^{\mathcal{A}}\right)^{k+j-q-1} \left(\tilde{f}_0^{\mathcal{A}}\right)^{k-q+p-1} \overline{b}^{\,l,i}_{(j,k-1)}.$$

In a similar way, we have

if the action of $E_{\mathcal{A}} b$ is given by (a1), then the action of $F_{\mathcal{A}}\left(\mathrm{C}_{\mathcal{A}}(b)\right)$ is given by (a2)′,

if the action of $E_{\mathcal{A}} b$ is given by (a2), then the action of $F_{\mathcal{A}}\left(\mathrm{C}_{\mathcal{A}}(b)\right)$ is given by (a1)′,

if the action of $E_{\mathcal{A}} b$ is given by (a3), then the action of $F_{\mathcal{A}}\left(\mathrm{C}_{\mathcal{A}}(b)\right)$ is given by (a5)′,

if the action of $E_{\mathcal{A}} b$ is given by (a4), then the action of $F_{\mathcal{A}}\left(\mathrm{C}_{\mathcal{A}}(b)\right)$ is given by (a4)′,

if the action of $E_{\mathcal{A}} b$ is given by (a5), then the action of $F_{\mathcal{A}}\left(\mathrm{C}_{\mathcal{A}}(b)\right)$ is given by (a3)′,

if the action of $E_{\mathcal{A}} b$ is given by (a6), then the action of $F_{\mathcal{A}}\left(\mathrm{C}_{\mathcal{A}}(b)\right)$ is given by (a6)′.

Thus we have $\mathrm{C}_{\mathcal{A}}\left(E_{\mathcal{A}}(b)\right) = F_{\mathcal{A}}\left(\mathrm{C}_{\mathcal{A}}(b)\right)$. □

For $a \in \mathbf{Z}$, we define a_+ by

$$a_+ = \begin{cases} a & (a > 0), \\ 0 & (a \le 0). \end{cases}$$

We set B_C, B_W, B_U, $B_R \subset \mathcal{A}^{(l)}$ by

$$B_C = \left\{ \left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{\,l,i}_{(l-i,j)} \;\middle|\; 0 \le i \le \left[\frac{l}{2}\right],\ i \le j \le l-i, 0 \le q \le p \le j \right\},$$

$$B_W = \left\{ \left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{\,l,i}_{(l-i,j)} \;\middle|\; 0 \le i \le \left[\frac{l}{2}\right],\ j = i,\ 0 \le p \le j,\ p < q \le y + j \right\},$$

$$B_U = \left\{ \left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{\,l,i}_{(l-i,j)} \;\middle|\; 0 \le i \le \left[\frac{l}{2}\right],\ i \le j \le l-i,\ p = j,\ j < q \le y + 2j - i \right\},$$

$$B_R = \left\{ \left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{\,l,i}_{(l-i,j)} \;\middle|\; \begin{array}{l} 0 \le i \le \left[\frac{l}{2}\right],\ i < j \le l-i, \\ 0 \le q < j,\ p < q \le y + j - (i-p)_+ \end{array} \right\},$$

where $y = y\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{\,l,i}_{(l-i,j)}\right) = \left[\frac{l-i-j}{3}\right]$.

In view of Definition 5.12, the conditions $\varepsilon_0^{\mathcal{A}}(b) = 0$, $\varphi_{\mathcal{A}}(b) = 0$ on $b = \left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{\,l,i}_{(l-i,j)}$ $(0 \le i \le \left[\frac{l}{2}\right], i \le j \le l-i, 0 \le p \le j)$ are rephrased as

$$\begin{cases} 0 \le q \le p - \left[\frac{-l+i+j}{3}\right] + (j-i) & (p \le i,\ l-i-j \equiv 0 \pmod 3), \\ 0 \le q \le p - 1 - \left[\frac{-l+i+j}{3}\right] + (j-i) & (p \le i,\ l-i-j \not\equiv 0 \pmod 3), \\ 0 \le q \le j - \left[\frac{-l+i+j}{3}\right] & (p > i,\ l-i-j \equiv 0 \pmod 3), \\ 0 \le q \le j - 1 - \left[\frac{-l+i+j}{3}\right] & (p > i,\ l-i-j \not\equiv 0 \pmod 3). \end{cases}$$

Summarizing, we have

$$0 \leq q \leq \left[\frac{l-i-j}{3}\right] + j - (i-p)_+ = y + j - (i-p)_+.$$

Thus we have

$$\begin{aligned}&\left\{b \in \mathcal{A} \,\middle|\, \varepsilon_0^{\mathcal{A}}(b) = 0, \varphi_{\mathcal{A}}(b) = 0\right\} \\ &= \left\{ \left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}_{(l-i,j)}^{l,i} \,\middle|\, 0 \leq i \leq \left[\frac{l}{2}\right], i \leq j \leq l-i, 0 \leq p \leq j, p \leq q \leq y + j - (i-p)_+ \right\}.\end{aligned}$$

Hence, we see easily that

$$\left\{b \in \mathcal{A} \,\middle|\, \varepsilon_0^{\mathcal{A}}(b) = 0, \varphi_{\mathcal{A}}(b) = 0\right\} \subset B_C \cup B_W \cup B_U \cup B_R.$$

By Proposition 5.3, we have

$$\left\{ \left(\tilde{f}_0^{\mathcal{A}}\right)^r \left(\tilde{f}_1^{\mathcal{A}}\right)^p \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}_{(l-i,j)}^{l,i} \,\middle|\, 0 \leq p \leq j, 0 \leq r \leq j-p \right\} = B_C. \tag{5.2.18}$$

Proposition 5.14 *For $b \in B_W \cup B_U$, we have*

$$\mathrm{C}_{\mathcal{A}}\left(E_{\mathcal{A}}^{\varepsilon_{\mathcal{A}}(b)}(b)\right) \in B_C.$$

Proof By calculation for $b = \left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}_{(l-i,i)}^{l,i} \in B_W$ $(0 \leq p < i, p < q \leq y + i)$ we have

$$\mathrm{C}_{\mathcal{A}}\left(E_{\mathcal{A}}^{\varepsilon_{\mathcal{A}}(b)} b\right) = \left(\tilde{f}_1^{\mathcal{A}}\right)^{i+2q-2p} \left(\tilde{f}_0^{\mathcal{A}}\right)^{2i+3q-4p} \overline{b}_{(l-p,2i+3q-4p)}^{l,p}.$$

Since $p < q$, $p < i$, we have

$$2i + 3q - 4p - (i + 2q - 2p) = (i-p) + (q-p) > 0.$$

Then we have

$$\mathrm{C}_{\mathcal{A}}\left(E_{\mathcal{A}}^{\varepsilon_{\mathcal{A}}(b)} b\right) \in B_C.$$

In a similar way, for $b \in B_U$ we have

$$\mathrm{C}_{\mathcal{A}}\left(E_{\mathcal{A}}^{\varepsilon_{\mathcal{A}}(b)} b\right) \in B_C.$$

□

Similar to Proposition 5.14, we have following proposition:

Proposition 5.15 *For $b \in B_R$, we have*

$$\mathrm{C}_{\mathcal{A}}\left(E_{\mathcal{A}}^{\varepsilon_{\mathcal{A}}(b)}(b)\right) \in B_R.$$

5.2.4 Definition of Φ on $\mathcal{A}$ and $\tilde{f}_0^{\mathcal{G}}$ on $\mathcal{G}^l$

In this section, we define one-to-one map $\Phi : \mathcal{A} \to \mathcal{G}^l$ and Kashiwara operator $\tilde{f}_0^{\mathcal{G}}$ on $\mathcal{G}^l$ expoiting operators on $\mathcal{A}$.

Since roots α_0 and α_2 of algebra $U_q(G_2^{(1)})$ are orthogonal, the Kashiwara operator $\tilde{f}_0^{\mathcal{G}}$ commutes with $\tilde{f}_2^{\mathcal{G}}$.

(D1) $\tilde{f}_0^{\mathcal{G}} \tilde{f}_2^{\mathcal{G}}(b) = \tilde{f}_2^{\mathcal{G}} \tilde{f}_0^{\mathcal{G}}(b)$, $\tilde{e}_0^{\mathcal{G}} \tilde{e}_2^{\mathcal{G}}(b) = \tilde{e}_2^{\mathcal{G}} \tilde{e}_0^{\mathcal{G}}(b)$, for $b \in \mathcal{G}^l$.

In view of Proposition 5.8, we are going to construct the one-to-one map

$$\Phi : \mathcal{A} \to \mathcal{G}^l,$$

with the following properties;

(E1) $\tilde{e}_1^{\mathcal{G}}\Phi(b)=\Phi\left(\tilde{e}_1^{\mathcal{A}}b\right)$, for $b\in\mathcal{A}$,

(E2) $\tilde{f}_2^{\mathcal{G}}\Phi(b)=\Phi\left(F_{\mathcal{A}}b\right)$, for $b\in\mathcal{A}$,

(E3) $\mathrm{wt}_1^{\mathcal{G}}\left(\Phi(b)\right)=\mathrm{wt}_1^{\mathcal{A}}(b)$, for $b\in\mathcal{A}$,

(E4) $\mathrm{wt}_2^{\mathcal{G}}\left(\Phi(b)\right)=-2\mathrm{wt}_1^{\mathcal{A}}(b)-\mathrm{wt}_0^{\mathcal{A}}(b)$, for $b\in\mathcal{A}$,

(E5) $\tilde{e}_0^{\mathcal{G}}\Phi(b)=0$ (resp. $\tilde{f}_0^{\mathcal{G}}\Phi(b)=0$) if and only if $\tilde{e}_0^{\mathcal{A}}(b)=0$ (resp. $\tilde{f}_0^{\mathcal{A}}(b)=0$), for $b\in\mathcal{A}$.

By (E1) and (E2), we see easily

(E1)$'$ $\tilde{f}_1^{\mathcal{G}}\Phi(b)=\Phi\left(\tilde{f}_1^{\mathcal{A}}b\right)$, for $b\in\mathcal{A}$,

(E2)$'$ $\tilde{e}_2^{\mathcal{G}}\Phi(b)=\Phi\left(E_{\mathcal{A}}b\right)$, for $b\in\mathcal{A}$.

We call an element $b\in\mathcal{A}$ $\mathcal{A}$*-terminal* if $F_{\mathcal{A}}(b)=0$. An $E_{\mathcal{A}}$*-string* is a sequence of elements in $\mathcal{A}$, b , $E_{\mathcal{A}}(b)$, $E_{\mathcal{A}}^2(b)$, …, $E_{\mathcal{A}}^n(b)$, where b is $\mathcal{A}$-terminal and $E_{\mathcal{A}}^{n+1}(b)=0$ ($n\in\mathbf{Z}_{\geq 0}$). We first define Φ for $\mathcal{A}$-terminal elements using (E3)(E4) mainly. Next we define the the action of Φ for elements of an $E_{\mathcal{A}}$-string using (E2). We verify (E1) completely in §5.3.

Definition 5.16 *We define*

$$\Phi\left(\left(\tilde{f}_0^{\mathcal{A}}\right)^p\overline{b}^{\,l,0}_{(l,l)}\right) = \left\llbracket\llbracket 6\rrbracket^p\llbracket\overline{2}\rrbracket^{l-p}\right\rrbracket \quad (0\leq p\leq l), \tag{5.2.19}$$

$$\Phi\left(\left(\tilde{e}_0^{\mathcal{A}}\right)^p\underline{b}^{\,l,0}_{(-l,-l)}\right) = \left\llbracket\llbracket 2\rrbracket^{l-p}\llbracket\overline{6}\rrbracket^p\right\rrbracket \quad (0\leq p\leq l). \tag{5.2.20}$$

We verify the properties (E1)–(E5).

By the definition of $\overline{b}^{\,l,0}_{(l,l)}$,

$$\mathrm{wt}^{\mathcal{A}}\left(\overline{b}^{\,l,0}_{(l,l)}\right)=l\Lambda_1+l\Lambda_0.$$

By (E3) and (E4), we see that $\Phi\left(\overline{b}^{\,l,0}_{(l,l)}\right)$ is the element with weight $l\Lambda_1-3l\Lambda_2$. We can see easily that $b=\left\llbracket\llbracket 6\rrbracket^k\llbracket\overline{2}\rrbracket^{l-k}\right\rrbracket$ $(0\leq k\leq l)$ is the only element such that $\mathrm{wt}_2^{\mathcal{G}}(b)=-3l$. Moreover $b=\left\llbracket\llbracket\overline{2}\rrbracket^l\right\rrbracket$ is the only element such that $\mathrm{wt}_1^{\mathcal{G}}(b)=l$. The highest weight element is only element in $B^i_{(k,j)}$ that satisfies $\mathrm{wt}(b)=k'\Lambda_1+j'\Lambda_0$ $(k'\geq k, j'\geq j)$. By the definition of $\mathcal{A}$, the element $b\in\mathcal{A}$ with $\mathrm{wt}(b)=l\Lambda_1+l\Lambda_0$ is $\overline{b}^{\,l,0}_{(l,l)}$. Therefore, we have

$$\Phi\left(\overline{b}^{\,l,0}_{(l,l)}\right)=\left\llbracket\llbracket\overline{2}\rrbracket^l\right\rrbracket. \tag{5.2.21}$$

We verify (E1). Since $\tilde{e}_1^{\mathcal{A}}\overline{b}^{\,l,0}_{(l,l)}=0$, we verify $\tilde{e}_1^{\mathcal{G}}\left\llbracket\llbracket\overline{2}\rrbracket^l\right\rrbracket=0$. Using Proposition 3.9,

$$\Psi_1\left(\left\llbracket\llbracket\overline{2}\rrbracket^l\right\rrbracket\right)=u_+^l.$$

Then we have

$$\tilde{e}_1^{\mathcal{G}}\left\llbracket\llbracket\overline{2}\rrbracket^l\right\rrbracket=0$$

In a similar way, we verify (E2). We consider $\left(\tilde{f}_0^{\mathcal{A}}\right)^p\overline{b}^{\,l,0}_{(l,l)}$. We see that the element $b\in\mathcal{G}^l$ which satisfy following formula is unique:

$$\mathrm{wt}^{\mathcal{G}}(b)=(l+k)\Lambda_1-3l\Lambda_2.$$

Thus, we can define

$$\Phi\left(\left(\tilde{f}_0^{\mathcal{A}}\right)^p\overline{b}^{\,l,0}_{(l,l)}\right)=\left\llbracket\llbracket 6\rrbracket^p\llbracket\overline{2}\rrbracket^{l-p}\right\rrbracket \quad (0\leq p\leq l). \tag{5.2.22}$$

Similarly, by the calculation of weight, we can define uniquely

$$\Phi\left(\left(\tilde{e}_0^{\mathcal{A}}\right)^p \underline{b}^{l,0}_{(-l,-l)}\right) = \left\llbracket \llbracket 2 \rrbracket^{l-p} \llbracket \overline{6} \rrbracket^{p} \right\rrbracket \quad (0 \le p \le l).$$

□

By (E5) and (5.2.22), we define the action of $\tilde{e}_0^{\mathcal{G}}$ and $\tilde{f}_0^{\mathcal{G}}$ for $\left\llbracket \llbracket 6 \rrbracket^{p} \llbracket \overline{2} \rrbracket^{l-p} \right\rrbracket$ $(0 \le p \le l)$ by

$$\tilde{f}_0^{\mathcal{G}} \left\llbracket \llbracket 6 \rrbracket^{p} \llbracket \overline{2} \rrbracket^{l-p} \right\rrbracket = \begin{cases} \left\llbracket \llbracket 6 \rrbracket^{p+1} \llbracket \overline{2} \rrbracket^{l-p-1} \right\rrbracket & 0 \le p \le l-1 \\ 0 & p = l \end{cases} \tag{5.2.23}$$

$$\tilde{e}_0^{\mathcal{G}} \left\llbracket \llbracket 6 \rrbracket^{p} \llbracket \overline{2} \rrbracket^{l-p} \right\rrbracket = \begin{cases} \left\llbracket \llbracket 6 \rrbracket^{p-1} \llbracket \overline{2} \rrbracket^{l-p+1} \right\rrbracket & 1 \le p \le l \\ 0 & p = 0 \end{cases} \tag{5.2.24}$$

In view of Definition 5.12, 5.16 and (E2)′, following definition is led.

Definition 5.17 *We define*

$$\Phi\left(\left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{l,0}_{(k,l)}\right) = \left(\tilde{e}_2^{\mathcal{G}}\right)^{l-k} \left\llbracket \llbracket 6 \rrbracket^{p} \llbracket \overline{2} \rrbracket^{l-p} \right\rrbracket. \tag{5.2.25}$$

where $0 \le k, p \le l$.

We verify the properties (E1)–(E5). The property (E2) is obvious. We verify (E1)(E3). We see

$$\varepsilon_1^{\mathcal{A}}\left(\left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{l,0}_{(k,l)}\right) = 0,$$

$$\varphi_1^{\mathcal{A}}\left(\left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{l,0}_{(k,l)}\right) = p + k.$$

On the other hand,

$$\begin{aligned} &\left(\tilde{e}_2^{\mathcal{G}}\right)^{l-k} \left\llbracket \llbracket 6 \rrbracket^{p} \llbracket \overline{2} \rrbracket^{l-p} \right\rrbracket \\ &\quad = \begin{cases} \left\llbracket \llbracket 2 \rrbracket^{(l-k)/3} \llbracket 6 \rrbracket^{p-(l-k)/3} \llbracket \overline{2} \rrbracket^{l-p} \right\rrbracket & 0 \le l-k \le 3p,\ \ l-k \equiv 0 \pmod 3 \\ \left\llbracket \llbracket 2 \rrbracket^{[(l-k)/3]} \llbracket 4 \rrbracket \llbracket 6 \rrbracket^{p-1-[(l-k)/3]} \llbracket \overline{2} \rrbracket^{l-p} \right\rrbracket & 0 < l-k < 3p,\ \ l-k \equiv 1 \pmod 3 \\ \left\llbracket \llbracket 2 \rrbracket^{[(l-k)/3]} \llbracket 3 \rrbracket \llbracket 6 \rrbracket^{p-1-[(l-k)/3]} \llbracket \overline{2} \rrbracket^{l-p} \right\rrbracket & 0 < l-k < 3p,\ \ l-k \equiv 2 \pmod 3 \\ \left\llbracket \llbracket 2 \rrbracket^{p} \llbracket \overline{\mathcal{W}} \rrbracket^{l-k-3p} \llbracket \overline{2} \rrbracket^{k+2p} \right\rrbracket & l-k > 3p \end{cases} \end{aligned}$$

Using Proposition3.9, we have

$$\varepsilon_1\left(\left(\tilde{e}_2^{\mathcal{G}}\right)^{l-k} \left\llbracket \llbracket 6 \rrbracket^{p} \llbracket \overline{2} \rrbracket^{l-p} \right\rrbracket\right) = 0, \tag{5.2.26}$$

$$\varphi_1\left(\left(\tilde{e}_2^{\mathcal{G}}\right)^{l-k} \left\llbracket \llbracket 6 \rrbracket^{p} \llbracket \overline{2} \rrbracket^{l-p} \right\rrbracket\right) = p + k. \tag{5.2.27}$$

By calculation, we can verify (E4). We verify (E5). By (5.2.24) and (D1), we must have

$$\tilde{e}_0^{\mathcal{G}}\left(\left(\tilde{e}_2^{\mathcal{G}}\right)^{l-k} \left\llbracket \llbracket \overline{2} \rrbracket^{l} \right\rrbracket\right) = 0 \tag{5.2.28}$$

□

By Definition 5.17 and (E1)′, following definition is led.

Definition 5.18 *We define*

$$\Phi\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^q\left(\tilde{f}_0^{\mathcal{A}}\right)^l\overline{b}_{(k,l)}^{\,l,0}\right)=\left(\tilde{f}_1^{\mathcal{G}}\right)^q\left(\Phi\left(\left(\tilde{f}_0^{\mathcal{A}}\right)^l\overline{b}_{(k,l)}^{\,l,0}\right)\right)\ \ (0\le p\le l+j).$$

We can verify (E1),(E2), (E3), (E4). We see easily

$$\tilde{f}_0^{\mathcal{A}}\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^q\left(\tilde{f}_0^{\mathcal{A}}\right)^l\overline{b}_{(k,l)}^{\,l,0}\right)=0\ \ (0\le q\le l).$$

By (5.1.11), we see

$$\tilde{e}_0^{\mathcal{A}}\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^q\left(\tilde{f}_0^{\mathcal{A}}\right)^l\overline{b}_{(k,l)}^{\,l,0}\right)=0\ \ (l\le q\le l+j).$$

In order to satisfy (E4), we define

$$\tilde{f}_0^{\mathcal{G}}\left(\Phi\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^q\left(\tilde{f}_0^{\mathcal{A}}\right)^l\overline{b}_{(k,l)}^{\,l,0}\right)\right)=0\ \ (0\le q\le l),$$

$$\tilde{e}_0^{\mathcal{G}}\left(\Phi\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^q\left(\tilde{f}_0^{\mathcal{A}}\right)^l\overline{b}_{(k,l)}^{\,l,0}\right)\right)=0\ \ (l\le q\le l+j).$$

For $b\in B_{(k,j)}^i$ we define $y(b)\in\mathbf{Z}_{\ge 0}$ by

$$y(b)=\left[\frac{l-i-j}{3}\right].$$

We often write y instead of $y(b)$ for simplicity.

Definition 5.19 *For $\overline{b}_{(l-i,j)}^{\,l,i}$, we define*

$$\Phi\left(\overline{b}_{(l-i,j)}^{\,l,i}\right)=\begin{cases}\left\llbracket \llbracket 6\rrbracket^{y}\llbracket \mathcal{C}\rrbracket^{y+i}\llbracket \overline{2}\rrbracket^{y+i}\llbracket \overline{2}\rrbracket^{j-i}\right\rrbracket & l-i-j\equiv 0\pmod 3,\\ \left\llbracket \llbracket 6\rrbracket^{y+1}\llbracket \mathcal{C}\rrbracket^{y+i-1}\llbracket 4\rrbracket\llbracket \overline{2}\rrbracket^{y+i}\llbracket \overline{2}\rrbracket^{j-i}\right\rrbracket & \begin{pmatrix} l-i-j\equiv 1\pmod 3\\ y+i>0\end{pmatrix},\\ \left\llbracket \llbracket \overline{5}\rrbracket\llbracket \overline{2}\rrbracket^{j-i}\right\rrbracket & \begin{pmatrix} l-i-j\equiv 1\pmod 3\\ y+i=0\end{pmatrix},\\ \left\llbracket \llbracket 6\rrbracket^{y+1}\llbracket \mathcal{C}\rrbracket^{y+i}\llbracket \overline{3}\rrbracket\llbracket \overline{2}\rrbracket^{y+i}\llbracket \overline{2}\rrbracket^{j-i}\right\rrbracket & (l-i-j\equiv 2\pmod 3).\end{cases}\tag{5.2.29}$$

We verify the properties (E1)–(E5). We verify (E1) and (E3). We see easily

$$\tilde{e}_1^{\mathcal{A}}\left(\overline{b}_{(l-i,j)}^{\,l,i}\right)=0,\ \tilde{f}_1^{\mathcal{G}}\left(\overline{b}_{(l-i,j)}^{\,l,i}\right)=l-i.$$

By proposition 3.9, we have

$$\begin{aligned}\Psi_1\left(\left\llbracket \llbracket 6\rrbracket^{y}\llbracket \mathcal{C}\rrbracket^{y+i}\llbracket \overline{2}\rrbracket^{y+j}\right\rrbracket\right)&=u_+^{y+j}u_-^{y+i}u_+^{y+i}u_+^{2y},\\ \mathrm{Red}_1\left(\left\llbracket \llbracket 6\rrbracket^{y}\llbracket \mathcal{C}\rrbracket^{y+i}\llbracket \overline{2}\rrbracket^{y+j}\right\rrbracket\right)&=u_+^{l-i}.\end{aligned}$$

Then we have

$$\tilde{e}_1^{\mathcal{G}}\left(\left\llbracket \llbracket 6\rrbracket^{y}\llbracket \mathcal{C}\rrbracket^{y+i}\llbracket \overline{2}\rrbracket^{y+j}\right\rrbracket\right)=0,$$

$$\tilde{f}_1^{\mathcal{G}}\left(\left\llbracket \llbracket 6\rrbracket^{y}\llbracket \mathcal{C}\rrbracket^{y+i}\llbracket \overline{2}\rrbracket^{y+j}\right\rrbracket\right)=l-i.$$

In a similar way, we have

$$\begin{aligned}
&\tilde{e}_1^{\mathcal{G}}\left(\left\llbracket \llbracket 6\rrbracket^{y+1}\llbracket \mathcal{C}\rrbracket^{y+i-1}\llbracket \overline{4}\rrbracket\llbracket \overline{2}\rrbracket^{y+j}\right\rrbracket\right)=0, && \tilde{f}_1^{\mathcal{G}}\left(\left\llbracket \llbracket 6\rrbracket^{y+1}\llbracket \mathcal{C}\rrbracket^{y+i-1}\llbracket \overline{4}\rrbracket\llbracket \overline{2}\rrbracket^{y+j}\right\rrbracket\right)=l-i,\\
&\tilde{e}_1^{\mathcal{G}}\left(\left\llbracket \llbracket \overline{5}\rrbracket\llbracket \overline{2}\rrbracket^{j-i}\right\rrbracket\right)=0, && \tilde{f}_1^{\mathcal{G}}\left(\left\llbracket \llbracket \overline{5}\rrbracket\llbracket \overline{2}\rrbracket^{j-i}\right\rrbracket\right)=l-i,\\
&\tilde{e}_1^{\mathcal{G}}\left(\left\llbracket \llbracket 6\rrbracket^{y+1}\llbracket \mathcal{C}\rrbracket^{y+i}\llbracket \overline{3}\rrbracket\llbracket \overline{2}\rrbracket^{y+j}\right\rrbracket\right)=0, && \tilde{e}_1^{\mathcal{G}}\left(\left\llbracket \llbracket 6\rrbracket^{y+1}\llbracket \mathcal{C}\rrbracket^{y+i}\llbracket \overline{3}\rrbracket\llbracket \overline{2}\rrbracket^{y+j}\right\rrbracket\right)=l-i,
\end{aligned}$$

By calculation, we can verify (E2) and (E4). We verify (E5). We put $b=\left\llbracket \llbracket 6\rrbracket^{y}\llbracket \mathcal{C}\rrbracket^{y+i}\llbracket \overline{2}\rrbracket^{y+j}\right\rrbracket$. By calculation and Lemma 5.17, we have

$$\begin{aligned}
\left(\tilde{e}_2^{\mathcal{G}}\right)^{2l-2i+j}b &= \left\llbracket \llbracket 2\rrbracket^{y(b)}\llbracket \mathcal{C}\rrbracket^{y(b)+i}\llbracket \overline{6}\rrbracket^{y(b)+j}\right\rrbracket,\\
\left(\tilde{e}_1^{\mathcal{G}}\right)^{l+j}\left(\tilde{e}_2^{\mathcal{G}}\right)^{2l-2i+j}b &= \left\llbracket \llbracket 2\rrbracket^{y(b)}\llbracket 6\rrbracket^{2y(b)+i+j}\right\rrbracket,\\
\left(\tilde{f}_2^{\mathcal{G}}\right)^{l-i-j}\left(\tilde{e}_1^{\mathcal{G}}\right)^{l+j}\left(\tilde{e}_2^{\mathcal{G}}\right)^{2l-2i+j}b &= \left\llbracket \llbracket 6\rrbracket^{l}\right\rrbracket.
\end{aligned}$$

By Definition 5.18, we must have

$$\Phi\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{l+j}\left(\tilde{f}_0^{\mathcal{A}}\right)^{l}\overline{b}^{\,l,0}_{(i+j,l)}\right)=\left\llbracket \llbracket 2\rrbracket^{y}\llbracket \mathcal{C}\rrbracket^{y+i}\llbracket \overline{6}\rrbracket^{y+j}\right\rrbracket,$$

$$\tilde{e}_0^{\mathcal{G}}\left\llbracket \llbracket 2\rrbracket^{y}\llbracket \mathcal{C}\rrbracket^{y+i}\llbracket \overline{6}\rrbracket^{y+j}\right\rrbracket=0.$$

By (D1), we must have

$$\tilde{e}_0^{\mathcal{G}}\left\llbracket \llbracket 6\rrbracket^{y}\llbracket \mathcal{C}\rrbracket^{y+i}\llbracket \overline{2}\rrbracket^{y+j}\right\rrbracket=0.$$

Similarly, we can prove other cases. □

Definition 5.20 *We set $b=\overline{b}^{\,l,i}_{(l-i,j)}$. We define the action of Φ as follows:*
If $0\le p\le i$,

$$\Phi\left(\left(\tilde{f}_0^{\mathcal{A}}\right)^{p}b\right)=\begin{cases}
\left\llbracket \llbracket 1\rrbracket^{p}\llbracket 6\rrbracket^{y}\llbracket \mathcal{C}\rrbracket^{y+i-p}\llbracket \overline{2}\rrbracket^{y+j}\right\rrbracket & (l-i-j\equiv 0 \pmod 3),\\
\left\llbracket \llbracket 1\rrbracket^{p}\llbracket 6\rrbracket^{y+1}\llbracket \mathcal{C}\rrbracket^{y+i-p-1}\llbracket \overline{4}\rrbracket\llbracket \overline{2}\rrbracket^{y+j}\right\rrbracket & (y+i>p,\ l-i-j\equiv 1 \pmod 3),\\
\left\llbracket \llbracket 1\rrbracket^{i}\llbracket \overline{5}\rrbracket\llbracket \overline{2}\rrbracket^{j}\right\rrbracket & (y+i=p,\ l-i-j\equiv 1 \pmod 3),\\
\left\llbracket \llbracket 1\rrbracket^{p}\llbracket 6\rrbracket^{y+1}\llbracket \mathcal{C}\rrbracket^{y+i-p}\llbracket \overline{3}\rrbracket\llbracket \overline{2}\rrbracket^{y+j}\right\rrbracket & (l-i-j\equiv 2 \pmod 3),
\end{cases}$$

if $i<p\le j$,

$$\Phi\left(\left(\tilde{f}_0^{\mathcal{A}}\right)^{p}b\right)=\begin{cases}
\left\llbracket \llbracket 1\rrbracket^{i}\llbracket 6\rrbracket^{p-i+y}\llbracket \mathcal{C}\rrbracket^{y}\llbracket \overline{2}\rrbracket^{y+j-p+i}\right\rrbracket & (l-i-j\equiv 0 \pmod 3),\\
\left\llbracket \llbracket 1\rrbracket^{i}\llbracket 6\rrbracket^{p-i+y+1}\llbracket \mathcal{C}\rrbracket^{y-1}\llbracket \overline{4}\rrbracket\llbracket \overline{2}\rrbracket^{y+j-p+i}\right\rrbracket & (y>0, l-i-j\equiv 1 \pmod 3),\\
\left\llbracket \llbracket 1\rrbracket^{i}\llbracket 6\rrbracket^{p-i}\llbracket \overline{5}\rrbracket\llbracket \overline{2}\rrbracket^{j-p+i}\right\rrbracket & (y=0, l-i-j\equiv 1 \pmod 3),\\
\left\llbracket \llbracket 1\rrbracket^{i}\llbracket 6\rrbracket^{p-i+y+1}\llbracket \mathcal{C}\rrbracket^{y}\llbracket \overline{3}\rrbracket\llbracket \overline{2}\rrbracket^{y+j-p+i}\right\rrbracket & (l-i-j\equiv 2 \pmod 3).
\end{cases}$$

By calculation, we can verify (E1)–(E4). In order to satisfy (E5), we define $\tilde{e}_0^{\mathcal{G}}$ on $\Phi\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^p \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}_{(l-i,j)}^{l,i}\right)$ $(0 \leq p \leq j)$ and $\tilde{f}_0^{\mathcal{G}}$ on $\Phi\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^j \overline{b}_{(l-i,j)}^{l,i}\right)$ $(0 \leq q \leq j)$.

Using Proposition 5.14, 5.15 and (E2), it is enough to define the action of Φ on $B_C \cup B_R$.

By (E1)$'$, we are led to the following definition.

Definition 5.21 *We define*

$$\Phi\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}_{(l-i,j)}^{l,i}\right) = \left(\tilde{f}_1^{\mathcal{G}}\right)^q \left(\Phi\left(\left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}_{(l-i,j)}^{l,i}\right)\right),$$

where $(0 \leq p \leq j,\ 0 \leq q \leq y + j - (i-p)_+)$.

Thus we have defined $\Phi(b)$ where $b \in B_C \cup B_R$.

Definition 5.22 *For* $b = \left\llbracket \llbracket b_1 \rrbracket \cdots \llbracket b_n \rrbracket \right\rrbracket \in B^{G_2}(n\Lambda_1)$ $(0 \leq n \leq l)$, *we define the involution on* $\mathcal{G}^l$ *by*

$$\mathrm{C}_{\mathcal{G}}(b) = \left\llbracket \llbracket \overline{b_n} \rrbracket \cdots \llbracket \overline{b_1} \rrbracket \right\rrbracket,$$

where if $b_i = \overline{m}$ $(m = 1, \ldots, 6)$ *then* $\llbracket \overline{b_i} \rrbracket = \llbracket m \rrbracket$, *if* $b_i = 0_1, 0_2$, *then* $\llbracket \overline{b_i} \rrbracket = \llbracket b_i \rrbracket$.

Remark 5.23 *Let* $\overline{b}$ *(resp.* $\underline{b}$*) be the highest (resp. lowest) weight element of* $B^{G_2}(l\Lambda_1)$*. By Proposition 3.9, 3.14 and 3.15, we have*

$$\overline{b} = \left\llbracket \llbracket 1 \rrbracket^l \right\rrbracket,$$

$$\underline{b} = \left\llbracket \llbracket \overline{1} \rrbracket^l \right\rrbracket.$$

By Defintion 5.22, we have

$$C_{\mathcal{G}}(\overline{b}) = \underline{b}.$$

Using Proposition 3.9 again, we see

$$\mathrm{C}_{\mathcal{G}}\left(\tilde{f}_{i_N} \cdots \tilde{f}_{i_1} \overline{b}\right) = \tilde{e}_{i_N} \cdots \tilde{e}_{i_1} \underline{b}.$$

Definition 5.24 *We define* Φ *for* $b \in B_C \cup B_R$

$$\Phi(E_{\mathcal{A}}^m b) = \left(\tilde{e}_2^{\mathcal{G}}\right)^m (\Phi(b)) \quad (0 \leq m \leq \varepsilon_{\mathcal{A}}(b)). \tag{5.2.30}$$

We define Φ *for* $b \in B_W \cup B_U$,

$$\Phi(E_{\mathcal{A}}^m b) = \mathrm{C}_{\mathcal{G}}\left(\Phi\left(\mathrm{C}_{\mathcal{A}}\left(E_{\mathcal{A}}^m b\right)\right)\right) \quad (0 \leq m \leq \varepsilon_{\mathcal{A}}(b)). \tag{5.2.31}$$

For $b \in B_R$, *we define*

$$\Phi\left(E_{\mathcal{A}}^m\left(\left(\tilde{f}_0^{\mathcal{A}}\right)^{\varphi_0(b)}(b)\right)\right) = \mathrm{C}_{\mathcal{G}}\left(\Phi\left(F_{\mathcal{A}}^{\varepsilon_{\mathcal{A}}(b)-m}\mathrm{C}_{\mathcal{A}}(b)\right)\right). \tag{5.2.32}$$

Remark 5.25 *By Definition 5.24, we see that the relation (5.2.30) satisfies* (E2)*. By (5.2.30) and Proposition 5.14, we see that* $\mathrm{C}_{\mathcal{A}}(E_{\mathcal{A}}^m b)$ *is already defined. By (5.2.31) ,We see* $\mathrm{C}_{\mathcal{A}}(\mathrm{C}_{\mathcal{A}}(E_{\mathcal{A}}^m b)) = b$ $(0 \leq m \leq \varepsilon_{\mathcal{A}}(b))$.

We have defined $\Phi(b)$ where $b \in \mathcal{A}^{(l)}$.

Proposition 5.26 *For $b \in \mathcal{A}^{(l)}$,*

$$\mathrm{C}_{\mathcal{G}}\left(\Phi(b)\right) = \Phi(\mathrm{C}_{\mathcal{A}}\left(b\right)).$$

Proof We put $b = \left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^j \overline{b}^{l,i}_{(l-i,j)}$. By (5.2.12), (5.2.13) and (5.2.14),

$$E_{\mathcal{A}}^{\varepsilon_{\mathcal{A}}(b)}\left(\left(\tilde{e}_0^{\mathcal{A}}\right)^{j-q} b\right) = \begin{cases} \left(\tilde{f}_1^{\mathcal{A}}\right)^{l+j-2q} \left(\tilde{f}_0^{\mathcal{A}}\right)^{l-q} \overline{b}^{l,q}_{(j+i-q,l-q)} & \text{if } 0 \le q \le \left[\dfrac{i+j}{2}\right], \\ \left(\tilde{f}_1^{\mathcal{A}}\right)^{l+j-2q} \left(\tilde{f}_0^{\mathcal{A}}\right)^{l-i-j+q} \overline{b}^{l,2i+2j-3q}_{(2i+2j-3q,l-2i-2j+3q)} & \text{if } \left[\dfrac{i+j}{2}\right] < q \le \dfrac{2j+i}{3}, \\ \left(\tilde{f}_1^{\mathcal{A}}\right)^{l-i-j+q} \left(\tilde{f}_0^{\mathcal{A}}\right)^{l-i-j+q} \overline{b}^{l,i}_{(-2j+3q,l-i)} & \text{if } \dfrac{2j+i}{3} < q \le j. \end{cases} \tag{5.2.33}$$

If $0 \le q \le \left[\frac{i+j}{2}\right]$,

$$\begin{aligned}
\Phi\left(\left(\tilde{f}_0^{\mathcal{A}}\right)^j \overline{b}^{l,i}_{(l-i,j)}\right) &= \left\llbracket \llbracket 1 \rrbracket^i \llbracket 6 \rrbracket^{j-i+y} \llbracket \mathcal{C} \rrbracket^y \llbracket \overline{2} \rrbracket^{y+i} \right\rrbracket, \\
\Phi\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^j \overline{b}^{l,i}_{(l-i,j)}\right) &= \begin{cases} \left\llbracket \llbracket 1 \rrbracket^i \llbracket 6 \rrbracket^{j-i+y} \llbracket \mathcal{C} \rrbracket^y \llbracket \overline{2} \rrbracket^{y+i-q} \llbracket \overline{1} \rrbracket^q \right\rrbracket & (i > q), \\ \left\llbracket \llbracket 1 \rrbracket^i \llbracket 6 \rrbracket^{j-q+y} \llbracket \mathcal{C} \rrbracket^{y+q-i} \llbracket \overline{2} \rrbracket^y \llbracket \overline{1} \rrbracket^i \right\rrbracket & (i \le q), \end{cases} \\
\Phi\left(E_{\mathcal{A}}^{\varepsilon_{\mathcal{A}}(b)} \left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^j \overline{b}^{l,i}_{(l-i,j)}\right) &= \begin{cases} \left\llbracket \llbracket 1 \rrbracket^i \llbracket 2 \rrbracket^{j-i+y} \llbracket \mathcal{C} \rrbracket^y \llbracket \overline{6} \rrbracket^{y+i-q} \llbracket \overline{1} \rrbracket^q \right\rrbracket & (i > q), \\ \left\llbracket \llbracket 1 \rrbracket^i \llbracket 2 \rrbracket^{j-q+y} \llbracket \mathcal{C} \rrbracket^{y+q-i} \llbracket \overline{6} \rrbracket^y \llbracket \overline{1} \rrbracket^i \right\rrbracket & (i \le q), \end{cases} \\
\mathrm{C}_{\mathcal{G}}\left(\Phi\left(E_{\mathcal{A}}^{\varepsilon_{\mathcal{A}}(b)} \left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^j \overline{b}^{l,i}_{(l-i,j)}\right)\right) &= \begin{cases} \left\llbracket \llbracket 1 \rrbracket^q \llbracket 6 \rrbracket^{i-q+y} \llbracket \mathcal{C} \rrbracket^y \llbracket \overline{2} \rrbracket^{y+j-i} \llbracket \overline{1} \rrbracket^i \right\rrbracket & (i > q), \\ \left\llbracket \llbracket 1 \rrbracket^i \llbracket 6 \rrbracket^{y} \llbracket \mathcal{C} \rrbracket^{y+q-i} \llbracket \overline{2} \rrbracket^{y+j-q} \llbracket \overline{1} \rrbracket^i \right\rrbracket & (i \le q). \end{cases}
\end{aligned}$$

On the other hand,

$$\begin{aligned}
\Phi\left(\left(\tilde{f}_0^{\mathcal{A}}\right)^i \overline{b}^{l,q}_{(l-q,i+j-q)}\right) &= \begin{cases} \left\llbracket \llbracket 1 \rrbracket^q \llbracket 6 \rrbracket^{i-q+y} \llbracket \mathcal{C} \rrbracket^y \llbracket \overline{2} \rrbracket^{y+j} \right\rrbracket & (i > q), \\ \left\llbracket \llbracket 1 \rrbracket^i \llbracket 6 \rrbracket^{y} \llbracket \mathcal{C} \rrbracket^{y+q-i} \llbracket \overline{2} \rrbracket^{y+j+i-q} \right\rrbracket & (i \le q), \end{cases} \\
\Phi\left(\mathrm{C}_{\mathcal{A}}\left(E_{\mathcal{A}}^{\varepsilon_{\mathcal{A}}(b)} b\right)\right) &= \Phi\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^i \left(\tilde{f}_0^{\mathcal{A}}\right)^i \overline{b}^{l,q}_{(l-q,i+j-q)}\right) \\
&= \begin{cases} \left\llbracket \llbracket 1 \rrbracket^q \llbracket 6 \rrbracket^{i-q+y} \llbracket \mathcal{C} \rrbracket^y \llbracket \overline{2} \rrbracket^{y+j-i} \llbracket \overline{1} \rrbracket^i \right\rrbracket & (i > q), \\ \left\llbracket \llbracket 1 \rrbracket^i \llbracket 6 \rrbracket^{y} \llbracket \mathcal{C} \rrbracket^{y+q-i} \llbracket \overline{2} \rrbracket^{y+j-q} \llbracket \overline{1} \rrbracket^i \right\rrbracket & (i \le q). \end{cases}
\end{aligned}$$

By (5.2.31), we have

$$\mathrm{C}_{\mathcal{G}}\left(\Phi\left(E_{\mathcal{A}}^m b\right)\right) = \Phi\left(\mathrm{C}_{\mathcal{A}}\left(E_{\mathcal{A}}^m b\right)\right) \quad (0 \le m \le \varepsilon_0(b))$$

In a similar way, we can calculate other cases. □

Definition 5.27 *For $b \in \mathcal{G}^l$, we define $\tilde{f}_0^{\mathcal{G}}(b)$ and $\tilde{e}_0^{\mathcal{G}}(b)$ by*

$$\tilde{f}_0^{\mathcal{G}}(b) = \Phi \tilde{f}_0^{\mathcal{A}} \Phi^{-1}(b),$$

$$\tilde{e}_0^{\mathcal{G}}(b) = \Phi\tilde{e}_0^{\mathcal{A}}\Phi^{-1}(b).$$

We define $\varepsilon_0^{\mathcal{G}}(b)$ and $\varphi_0^{\mathcal{G}}(b)$ by

$$\varepsilon_0^{\mathcal{G}}(b) = \max\left\{n \,\middle|\, \left(\tilde{e}_0^{\mathcal{G}}\right)^n b \neq 0\right\},$$

$$\varphi_0^{\mathcal{G}}(b) = \max\left\{n \,\middle|\, \left(\tilde{f}_0^{\mathcal{G}}\right)^n b \neq 0\right\}.$$

By (C2), it is obvious that $\tilde{f}_0^{\mathcal{G}} = \Phi\tilde{f}_0^{\mathcal{A}}\Phi^{-1}$, $\tilde{e}_0^{\mathcal{G}} = \Phi\tilde{e}_0^{\mathcal{A}}\Phi^{-1}$ satisfy (D1).

Remark 5.28 *By Difinition 5.27, for $b \in \mathcal{A}$ we have*

$$\varepsilon_0^{\mathcal{G}}(\Phi(b)) = \varepsilon_0^{\mathcal{A}}(b),$$

$$\varphi_0^{\mathcal{G}}(\Phi(b)) = \varphi_0^{\mathcal{A}}(b).$$

Proposition 5.29 *The action of $\tilde{f}_0^{\mathcal{G}}$ is unique.*

Proof Let Φ, Φ' be one-to-one maps $\mathcal{A} \to \mathcal{G}^l$ which satisfy conditions (E1)(E2)(E3) in §5.2.4. Here, by (E2) we have $\Phi F_{\mathcal{A}} = \tilde{f}_2^{\mathcal{G}}\Phi$, $\Phi' F_{\mathcal{A}} = \tilde{f}_2^{\mathcal{G}}\Phi'$. We see that if for $b \in \mathcal{A}$

$$\Phi(\tilde{f}_0^{\mathcal{A}}b) \neq \Phi'(\tilde{f}_0^{\mathcal{A}}b), \quad \text{for some } b \in \mathcal{A}$$

we have

$$\Phi\left(\tilde{f}_0^{\mathcal{A}}F_{\mathcal{A}}^{\varphi_{\mathcal{A}}(b)}(b)\right) \neq \Phi'\left(\tilde{f}_0^{\mathcal{A}}F_{\mathcal{A}}^{\varphi_{\mathcal{A}}(b)}(b)\right).$$

Therefore it is sufficient to verify that $\Phi(b)$ is unique, for $b \in \mathcal{A}$ such that $F_{\mathcal{A}}(b) = 0$. By (E1), $\Phi\tilde{f}_1^{\mathcal{A}} = \tilde{f}_1^{\mathcal{G}}\Phi$, then it is enough to verify that

$$\Phi\left(\left(\tilde{f}_0^{\mathcal{A}}\right)^p \bar{b}_{(l-i,j)}^{l,i}\right) \quad \left(0 \leq i \leq \left[\frac{l}{2}\right],\, 0 \leq j \leq l-i,\, 0 \leq p \leq j\right)$$

is unique. This is obvious by Proposition 5.20. □

The affine crystal B^l which is constructed with $\mathcal{G}$ and $\tilde{f}_0^{\mathcal{G}}$ satisfies Proposition 2.1.

5.3 Proof of commutativity of $\Phi\tilde{f}_1^{\mathcal{A}} = \tilde{f}_1^{\mathcal{G}}\Phi$

In this section we prove (E1) in §5.2.4, namely for $b \in \mathcal{A}$

$$\begin{tikzcd}
b \arrow[r] \arrow[d] & \Phi(b) \arrow[d] \\
\tilde{f}_1^{\mathcal{A}}b \arrow[r] & \tilde{f}_1^{\mathcal{G}}(\Phi(b)).
\end{tikzcd}$$

In the proof of commutativity we use $\tilde{e}_1^{\mathcal{A}}$, $\tilde{e}_1^{\mathcal{G}}$ instead of $\tilde{f}_1^{\mathcal{A}}$, $\tilde{f}_1^{\mathcal{G}}$ respectively. Consider an $E_{\mathcal{A}}$-string b, $E_{\mathcal{A}}(b)$, …, $E_{\mathcal{A}}^{\varphi_{\mathcal{A}}(b)}(b)$, where $b \in \mathcal{A}$ is $\mathcal{A}$-terminal. For any $b \in \mathcal{A}$, we can denote

$$b = E_{\mathcal{A}}^{n'}b',$$

where $n' \in \mathbf{Z}_{\geq 0}$, $b' \in \mathcal{A}$ is an $\mathcal{A}$-terminal element, namely $F_{\mathcal{A}}(b') = 0$. Similarly, we can denote

$$\tilde{e}_1^{\mathcal{A}}b = E_{\mathcal{A}}^{n''}b'',$$

where $n'' \in \mathbf{Z}_{\geq 0}$, $b'' \in \mathcal{A}$ is an A-terminal element. In order to show (E1), we will verify

$$\tilde{e}_1^{\mathcal{G}}\left(\Phi\left(E_{\mathcal{A}}^{n'}b'\right)\right) = \Phi\left(E_{\mathcal{A}}^{n''}b''\right). \tag{5.3.1}$$

Let us define $\mathcal{A}_+$, $\mathcal{A}_-$ by

$$\mathcal{A}_+ = \bigoplus_{i=0}^{\left[\frac{l}{2}\right]} \bigoplus_{i \le k \le j \le l-i} B^i_{(k,j)},$$

$$\mathcal{A}_- = \bigoplus_{i=0}^{\left[\frac{l}{2}\right]} \bigoplus_{i \le j < k \le l-i} B^i_{(k,j)}.$$

For $b' \in \mathcal{A}_+$ such that $E_{\mathcal{A}}(b') \in \mathcal{A}_-$, we see that the action of $E_{\mathcal{A}}$ is given by (a3) or (a4) or (a5). We take $b = \left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{l,i}_{(l-i,j)} \in B_C$ $(0 \le q \le p \le j)$, which is an $\mathcal{A}$-terminal element of an $E_{\mathcal{A}}$-string. By consulting the proof of (C3) in §5.2.2, we see that for b the action of $E_{\mathcal{A}}$ has following relations, using sequence depending b, $0 \le n_1 \le n_2 \le \cdots \le n_6 = \varepsilon_{\mathcal{A}}(b)$.
the action of $E_{\mathcal{A}}$ on $E^{n'}_{\mathcal{A}} b$ $(0 \le n' < n_1)$ is given by (a1), if $n_1 \ne 0$,
the action of $E_{\mathcal{A}}$ on $E^{n'}_{\mathcal{A}} b$ $(n_1 \le n' < n_2)$ is given by (a3), if $n_1 < n_2$,
the action of $E_{\mathcal{A}}$ on $E^{n'}_{\mathcal{A}} b$ $(n_2 \le n' < n_3)$ is given by (a4), if $n_2 < n_3$,
the action of $E_{\mathcal{A}}$ on $E^{n'}_{\mathcal{A}} b$ $(n_3 \le n' < n_4)$ is given by (a5), if $n_3 < n_4$,
the action of $E_{\mathcal{A}}$ on $E^{n'}_{\mathcal{A}} b$ $(n_4 \le n' < n_5)$ is given by (a2), if $n_4 < n_5$,
the action of $E_{\mathcal{A}}$ on $E^{n_6}_{\mathcal{A}} b$ is given by (a6).
Then we see that the number $\bar{n}(b) \in \mathbf{Z}_{\ge 0}$ which satisfy following condition is at most one:

$$E^{\bar{n}(b)}_{\mathcal{A}} b \in \mathcal{A}_+,$$

$$E^{\bar{n}(b)+1}_{\mathcal{A}} b \in \mathcal{A}_-.$$

If there does not exist such $\bar{n}(b)$, we put

$$\bar{n}(b) = \varepsilon_{\mathcal{A}}(b) = 2l - 2i + j - 3q.$$

By Proposition 5.13 and induction (5.2.6) it is sufficient to prove (5.3.1) for $b' = E^n_{\mathcal{A}} b$ $(0 \le n \le \bar{n}(b), b \in B_C)$.

5.3.1 Actions of $\tilde{e}_1^{\mathcal{A}}$

Proposition 5.30 *For* $\left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{l,i}_{(l-i,j)} \in B_C$ $(0 \le q \le p \le j)$, *we have following relations.*

Put $b_1 = \left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{l,i}_{(l-i,j)}$ $(3q \ge i + 2j - \left[\frac{j-i}{2}\right],\ p + 2q > i + 2j)$. *We have*

$$\tilde{e}_1^{\mathcal{A}}\left(E^n_{\mathcal{A}} b_1\right) = E^n_{\mathcal{A}}\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q-1}\left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{l,i}_{(l-i,j)}\right), \tag{5.3.2}$$

$$\tilde{e}_1^{\mathcal{A}}\left(E^{l-i-j+3(j-q)+1}_{\mathcal{A}}\ b_1\right) = E^{l-i-j+3(j-q)}_{\mathcal{A}}\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q}\left(\tilde{f}_0^{\mathcal{A}}\right)^{p+1} \overline{b}^{l,i}_{(l-i,j+1)}\right), \tag{5.3.3}$$

$$\tilde{e}_1^{\mathcal{A}}\left(E^{l-i-j+3(j-q)+2}_{\mathcal{A}}\ b_1\right) = E^{l-i-j+3(j-q)}_{\mathcal{A}}\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q+1}\left(\tilde{f}_0^{\mathcal{A}}\right)^{p+2} \overline{b}^{l,i}_{(l-i,j+2)}\right), \tag{5.3.4}$$

$$\tilde{e}_1^{\mathcal{A}}\left(E^{l-i-j+3(j-q+1)+n'}_{\mathcal{A}}\ b_1\right) = E^{l-i-j+3(j-q)+n'}_{\mathcal{A}}\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q+2}\left(\tilde{f}_0^{\mathcal{A}}\right)^{p+3} \overline{b}^{l,i}_{(l-i,j+3)}\right), \tag{5.3.5}$$

where $0 \le n \le l - j - i + 3(j-q)$, $0 \le n' \le 3(y-1)$, $y = \left[\frac{l-i-j}{3}\right]$.

Put $b_2 = \left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{l,i}_{(l-i,j)}$ $(p - q \le \left[\frac{j-i}{2}\right],\ p + 2q \le i + 2j)$. *We have*

$$\tilde{e}_1^{\mathcal{A}}\left(E^n_{\mathcal{A}}\ b_2\right) = E^n_{\mathcal{A}}\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q-1}\left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{l,i}_{(l-i,j)}\right), \tag{5.3.6}$$

$$\tilde{e}_1^{\mathcal{A}}\left(E^{l-2i+[(j-i)/2]-n'}_{\mathcal{A}}\ b_2\right) = E^{l-2i+[(j-i)/2]-3-n'}_{\mathcal{A}}\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q}\left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{l,i+1}_{(l-(i+1),j-1)}\right), \tag{5.3.7}$$

where $0 \le n \le l-2i+p-q$, $0 \le n' \le \left[\frac{j-i}{2}\right]$.

Put $b_3 = \left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{l,i}_{(l-i,j)}$ $(3q < i+2j-\left[\frac{j-i}{2}\right],\ \left[\frac{j-i}{2}\right] < p-q)$*, then we have*

$$\tilde{e}_1^{\mathcal{A}}\left(E_{\mathcal{A}}^n\, b_3\right) \quad = \quad E_{\mathcal{A}}^n\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q-1}\left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{l,i}_{(l-i,j)}\right), \tag{5.3.8}$$

where $0 \le n \le l-2i+\left[\frac{j-i}{2}\right]$.

Proof We consider the action of $\tilde{e}_1^{\mathcal{A}}$ for $\left(\tilde{f}_0^{\mathcal{A}}\right)^r \left(\tilde{f}_1^{\mathcal{A}}\right)^q \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{l,i}_{(k,j)}$ $(0 \le p \le j, p \le q \le p+k, 0 \le r \le j+q-2p)$. By Proposition 5.4, if $q=p$ we have

$$\tilde{e}_1^{\mathcal{A}}\left(\tilde{f}_0^{\mathcal{A}}\right)^r \left(\tilde{f}_1^{\mathcal{A}}\right)^p \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{l,i}_{(k,j)} = \left(\tilde{f}_0^{\mathcal{A}}\right)^r \left(\tilde{f}_1^{\mathcal{A}}\right)^{p-1} \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{l,i}_{(k,j)}$$

where $0 \le r \le j-p$. If $q > p$,

$$\tilde{e}_1^{\mathcal{A}}\left(\tilde{f}_0^{\mathcal{A}}\right)^r \left(\tilde{f}_1^{\mathcal{A}}\right)^p \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{l,i}_{(k,j)} \neq \left(\tilde{f}_0^{\mathcal{A}}\right)^r \left(\tilde{f}_1^{\mathcal{A}}\right)^{p-1} \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{l,i}_{(k,j)}$$

where $q-p \le r \le j+q-2p$.

We consider the case of $q=p$ in part I, and the case of $q>p$ in part II.

Part I We consider $\left(\tilde{f}_0^{\mathcal{A}}\right)^r \left(\tilde{f}_1^{\mathcal{A}}\right)^p \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{l,i}_{(k,j)}$ $(0 \le p \le j, 0 \le r \le j-p)$.

We define $b, b' \in B^i_{(k,j)}$ by

$$\begin{aligned} b &= \left(\tilde{f}_1^{\mathcal{A}}\right)^p \left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{l,i}_{(k,j)}, \\ b' &= \begin{cases} \left(\tilde{f}_1^{\mathcal{A}}\right)^{p-1} \left(\tilde{f}_0^{\mathcal{A}}\right)^{p-1} \overline{b}^{l,i}_{(k,j)} & (p>0) \\ 0 & (p=0) \end{cases}. \end{aligned}$$

Then by Proposition 5.3 we have

$$\tilde{e}_1^{\mathcal{A}}\left(\tilde{f}_0^{\mathcal{A}}\right)^r b = \left(\tilde{f}_0^{\mathcal{A}}\right)^{r+1} b',$$

(Figure 2). We define m_b by

$$m_b = j-k-3(j-p) = -2j-k+3p,$$

We consider an $E_{\mathcal{A}}$-string. We can express b using an $\mathcal{A}$-terminal element. By Definition 5.12, we have

$$b = E_{\mathcal{A}}^{l-i-k+(m_b)_+}\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{p-(m_b)_+}\left(\tilde{f}_0^{\mathcal{A}}\right)^p \overline{b}^{l,i}_{(l-i,j-(m_b)_+)}\right) \tag{5.3.9}$$

If $m_b < 0$ the action of $E_{\mathcal{A}}$ is given by (a1) or (a3) of Definition 5.12, and if $m_b \ge 0$ the action is given by (a2).

The case of (a1)**:**$E_{\mathcal{A}} b \in B^i_{(k-1,j)}$**.**

This case is $b = E_{\mathcal{A}}^{n'} b_1$ $(0 \le n' < \min\{l-2i, l-i+2j-3q\})$, $b = E_{\mathcal{A}}^{n'} b_2$ $(0 \le n' < l-2i)$, $b = E_{\mathcal{A}}^{n'} b_3$ $(0 \le n' < l-2i)$.
If $p=0$, we have

$$\tilde{e}_1^{\mathcal{A}} E_{\mathcal{A}} b = 0.$$


$\overline{b}^{l,i}_{(k,j)}$
p–1
p–1
j–p
b'
b
r
$\swarrow \tilde{f}^{\mathcal{A}}_1 \searrow \tilde{f}^{\mathcal{A}}_0$

Figure 2: Part I, the location of b and b' in the crystal graph of $B^i_{(k,j)}$.


$\overline{b}^{l,i}_{(k-1,j)}$
p–1
p–1
j–p
$E_{\mathcal{A}}b'$
$E_{\mathcal{A}}b$
r
$\swarrow \tilde{f}^{\mathcal{A}}_1 \searrow \tilde{f}^{\mathcal{A}}_0$

Figure 3: Part I, the case of (a1), the location of $E_{\mathcal{A}}b$ and $E_{\mathcal{A}}b'$ in the crystal graph of $B^i_{(k-1,j)}$.

If $p > 0$, since $m_b < 0$, we have $m_{b'} < 0$. Then the actions of $E_{\mathcal{A}}$ for b' is given by (a1). Then we have

$$\begin{aligned} E_{\mathcal{A}}b &= \left(\tilde{f}^{\mathcal{A}}_0\right)^p \left(\tilde{f}^{\mathcal{A}}_1\right)^p \overline{b}^{l,i}_{(k-1,j)}, \\ E_{\mathcal{A}}b' &= \left(\tilde{f}^{\mathcal{A}}_0\right)^{p-1} \left(\tilde{f}^{\mathcal{A}}_1\right)^{p-1} \overline{b}^{l,i}_{(k-1,j)}, \end{aligned}$$

(Figure 3). Then we obtain

$$\tilde{e}^{\mathcal{A}}_1 E_{\mathcal{A}}\left(\left(\tilde{f}^{\mathcal{A}}_0\right)^r b\right) = E_{\mathcal{A}}\left(\left(\tilde{f}^{\mathcal{A}}_0\right)^{r+1} b'\right). \tag{5.3.10}$$

We have the case of $(0 \leq n \leq \min\{l-2i, l-i+2j-3q\})$ of (5.3.2), the case of $(0 \leq n \leq l-2i)$ of (5.3.6), and the case of $(0 \leq n \leq l-2i)$ of (5.3.8),

The case of (a2)**:**$E_{\mathcal{A}}b \in B^i_{(k,j+1)}$**.**

This case is $b = E^{n'}_{\mathcal{A}} b_1$ $(l-i+2j-3q \leq n' < 2l-2i+j-3q, l-i+2j-3q < l-2i)$.
We denote

$$E_{\mathcal{A}}b = \left(\tilde{f}^{\mathcal{A}}_1\right)^{p+1} \left(\tilde{f}^{\mathcal{A}}_0\right)^{p+1} \overline{b}^{l,i}_{(k,j+1)}.$$

if $m_b = 0, 1, 2$, since $m_{b'} < 0 \leq m_b$, then the action of $E_{\mathcal{A}}$ for b' is given by (a1). Then we have

$$E_{\mathcal{A}}b' = \left(\tilde{f}^{\mathcal{A}}_1\right)^{p-1} \left(\tilde{f}^{\mathcal{A}}_0\right)^{p-1} \overline{b}^{l,i}_{(k-1,j)}.$$

Therefore, we have

$$\tilde{e}^{\mathcal{A}}_1 E_{\mathcal{A}}\left(\left(\tilde{f}^{\mathcal{A}}_0\right)^r b\right) \neq E_{\mathcal{A}}\left(\left(\tilde{f}^{\mathcal{A}}_0\right)^{r+1} b'\right).$$

By (5.3.9), we have

$$\tilde{e}_1^{\mathcal{A}} E_{\mathcal{A}} \left(\left(\tilde{f}_0^{\mathcal{A}} \right)^r \right) b = E_{\mathcal{A}}^{l-i-k} \left(\left(\tilde{f}_0^{\mathcal{A}} \right)^{r+1} \left(\tilde{f}_1^{\mathcal{A}} \right)^p \left(\tilde{f}_0^{\mathcal{A}} \right)^p \overline{b}^{l,i}_{(l-i,j+1)} \right).$$

if $m_b \geq 3$, then the action of $E_{\mathcal{A}}$ for b' is given by (a2), since $0 \leq m_{b'} < m_b$. Thus we have

$$E_{\mathcal{A}} b' = \left(\tilde{f}_1^{\mathcal{A}} \right)^p \left(\tilde{f}_0^{\mathcal{A}} \right)^p \overline{b}^{l,i}_{(k,j+1)},$$

(Figure 4). Therefore, we have

$$\begin{aligned} \tilde{e}_1^{\mathcal{A}} E_{\mathcal{A}} \left(\left(\tilde{f}_0^{\mathcal{A}} \right)^r b \right) &= E_{\mathcal{A}} \left(\left(\tilde{f}_0^{\mathcal{A}} \right)^{r+1} b' \right) \\ &= E_{\mathcal{A}}^{l-i-k+m_b-2} \left(\left(\tilde{f}_0^{\mathcal{A}} \right)^{r+1} \left(\tilde{f}_1^{\mathcal{A}} \right)^{p-m_b+2} \left(\tilde{f}_0^{\mathcal{A}} \right)^p \overline{b}^{l,i}_{(l-i,j-m_b+3)} \right). \end{aligned} \tag{5.3.11}$$

Therefore we have (5.3.3), (5.3.4), (5.3.5), if $l-i+2j-3q < l-2i$.

The case of (a3)**:**$E_{\mathcal{A}} b \in B^{i+1}_{(i+1,j-1).}$ **.**

This case is $b = E_{\mathcal{A}}^{l-2i} b_1$, $b = E_{\mathcal{A}}^{l-2i} b_2$, $b = E_{\mathcal{A}}^{l-2i} b_3$.
We denote

$$E_{\mathcal{A}} b = \left(\tilde{f}_1^{\mathcal{A}} \right)^{p+1} \left(\tilde{f}_0^{\mathcal{A}} \right)^p \overline{b}^{l,i+1}_{(k+1,j-1)},$$

Since $m_{b'} < m_b < 0$ the actions of $E_{\mathcal{A}}$ for b' is given by (a3). Then we have

$$E_{\mathcal{A}} b' = \left(\tilde{f}_1^{\mathcal{A}} \right)^p \left(\tilde{f}_0^{\mathcal{A}} \right)^{p-1} \overline{b}^{l,i+1}_{(k+1,j-1)}.$$

Here by Proposition 5.4 we have

$$\tilde{e}_1^{\mathcal{A}} \left(\tilde{f}_0^{\mathcal{A}} \right)^r \left(\tilde{f}_1^{\mathcal{A}} \right)^{p+1} \left(\tilde{f}_0^{\mathcal{A}} \right)^p \overline{b}^{l,i}_{(k,j)} = \left(\tilde{f}_0^{\mathcal{A}} \right)^{r+1} \left(\tilde{f}_1^{\mathcal{A}} \right)^p \left(\tilde{f}_0^{\mathcal{A}} \right)^{p-1} \overline{b}^{l,i}_{(k,j)} \neq \left(\tilde{f}_0^{\mathcal{A}} \right)^r \left(\tilde{f}_1^{\mathcal{A}} \right)^p \left(\tilde{f}_0^{\mathcal{A}} \right)^q \overline{b}^{l,i}_{(k,j)} \ (r > 0),$$

$$\tilde{e}_1^{\mathcal{A}} \left(\tilde{f}_1^{\mathcal{A}} \right)^{p+1} \left(\tilde{f}_0^{\mathcal{A}} \right)^p \overline{b}^{l,i}_{(k,j)} = \left(\tilde{f}_1^{\mathcal{A}} \right)^p \left(\tilde{f}_0^{\mathcal{A}} \right)^p \overline{b}^{l,i}_{(k,j)} \neq \tilde{f}_0^{\mathcal{A}} \left(\tilde{f}_1^{\mathcal{A}} \right)^p \left(\tilde{f}_0^{\mathcal{A}} \right)^{p-1} \overline{b}^{l,i}_{(k,j)},$$

(Figure 5). Then we have

$$\tilde{e}_1^{\mathcal{A}} \left(\tilde{f}_0^{\mathcal{A}} \right)^r E_{\mathcal{A}} b = \left(\tilde{f}_0^{\mathcal{A}} \right)^{r+1} E_{\mathcal{A}} b' \ \ (r > 0), \tag{5.3.12}$$

$$\tilde{e}_1^{\mathcal{A}} E_{\mathcal{A}} b = E_{\mathcal{A}}^{l-2(i+1)} \left(\left(\tilde{f}_1^{\mathcal{A}} \right)^p \left(\tilde{f}_0^{\mathcal{A}} \right)^p \overline{b}^{l,i+1}_{(l-i-1,j-1)} \right). \tag{5.3.13}$$

Therefore we have the case of $n = l-2i+1$ of (5.3.2), (5.3.6), (5.3.8).

Part II. We consider $\tilde{e}_1^{\mathcal{A}} E_{\mathcal{A}} \left(\tilde{f}_0^{\mathcal{A}} \right)^r \left(\tilde{f}_1^{\mathcal{A}} \right)^{p+s} \left(\tilde{f}_0^{\mathcal{A}} \right)^p \overline{b}^{l,i}_{(k,j)}$ $(i \leq k, j \leq l-i, \ k-i \geq s > 0)$
If $i < k, j \leq l-i$, we use induction (5.2.6). Otherwise, using involution, we have

$$\mathrm{C}_{\mathcal{A}} \left(\overline{b}^{l,i}_{(j,i)} \right) = \left(\tilde{f}_0^{\mathcal{A}} \right)^i \left(\tilde{f}_1^{\mathcal{A}} \right)^{i+j} \left(\tilde{f}_0^{\mathcal{A}} \right)^j \overline{b}^{l,i}_{(i,j)}.$$

Then it is sufficient to consider following element:

$$\tilde{e}_1^{\mathcal{A}} E_{\mathcal{A}} \left(\tilde{f}_0^{\mathcal{A}} \right)^r \left(\tilde{f}_1^{\mathcal{A}} \right)^{p+s} \left(\tilde{f}_0^{\mathcal{A}} \right)^p \overline{b}^{l,i}_{(i,j)} \ \ (i < j \leq l-i, \ 0 < s \leq k-i).$$


$\overline{b}^{l,i}_{(k,j+1)}$
p
p
j–p
$E_{\mathcal{A}}b$
r
$\swarrow \tilde{f}_1^{\mathcal{A}}$ $\searrow \tilde{f}_0^{\mathcal{A}}$

Figure 4: Part I, the case of (a2), the localtion of $E_{\mathcal{A}}b$ in the crystal graph of $B^i_{(k,j+1)}$.


$\overline{b}^{l,i+1}_{(k+1,j-1)}$
p–1
p
j–p–1
$E_{\mathcal{A}}b'$
$E_{\mathcal{A}}b$
j–p
$\swarrow \tilde{f}_1^{\mathcal{A}}$ $\searrow \tilde{f}_0^{\mathcal{A}}$

Figure 5: Part I, the case of (a3), the location of $E_{\mathcal{A}}b$ and $E_{\mathcal{A}}b'$ in the crystal graph $B^{i+1}_{(k+1,j-1)}$.

We denote $b, b', b'' \in B^i_{(k,j)}$ by

$$\begin{aligned} b &= \left(\tilde{f}_1^{\mathcal{A}}\right)^{p+s} \left(\tilde{f}_0^{\mathcal{A}}\right)^{p} \overline{b}^{l,i}_{(k,j)}, \\ b' &= \left(\tilde{f}_1^{\mathcal{A}}\right)^{p+s+1} \left(\tilde{f}_0^{\mathcal{A}}\right)^{p-1} \overline{b}^{l,i}_{(k,j)}, \\ b'' &= \tilde{e}_1^{\mathcal{A}} b. \end{aligned}$$

Then we have

$$\begin{aligned} \tilde{e}_1^{\mathcal{A}} \left(\tilde{f}_0^{\mathcal{A}}\right)^{r} b &= \left(\tilde{f}_0^{\mathcal{A}}\right)^{r} b'' \ (r < s), \\ \tilde{e}_1^{\mathcal{A}} \left(\tilde{f}_0^{\mathcal{A}}\right)^{r} b &= \left(\tilde{f}_0^{\mathcal{A}}\right)^{r+1} b' \ (r \geq s), \end{aligned}$$

(Figure 6). We rewrite m_b

$$m_b = (j+s) - (k-s) + s - 3((j+s) - p) = -2j - k + 3p.$$

Then we denote b using the $\mathcal{A}$-terminal element in an $E_{\mathcal{A}}$-string.

$$b = \begin{cases} E_{\mathcal{A}}^{s+l-2(k-s+1)} \left(\tilde{f}_1^{\mathcal{A}}\right)^{p} \left(\tilde{f}_0^{\mathcal{A}}\right)^{p} \overline{b}^{l,k-s+1}_{(l-(k-s+1),j+s)} & \text{if } m_b < 0, \\ E_{\mathcal{A}}^{s+l-2(k-s+1)+m_b} \left(\tilde{f}_1^{\mathcal{A}}\right)^{p-m_b} \left(\tilde{f}_0^{\mathcal{A}}\right)^{p-m_b} \overline{b}^{l,k-s+1}_{(l-(k-s+1),j+s-m_b)}, & \text{if } m_b \geq 0. \end{cases} \tag{5.3.14}$$

The case of (a2)**:** $E_{\mathcal{A}}b \in B^{i}_{(i,j+1)}$**.**

This case is $b = E_{\mathcal{A}}^{n'} b_1$ $(l-2i \leq l-i+2j-3q \leq n' < 2l-2i+j-3q)$.
We denote

$$E_{\mathcal{A}}b = \left(\tilde{f}_1^{\mathcal{A}}\right)^{p+s+1} \left(\tilde{f}_0^{\mathcal{A}}\right)^{p+1} \overline{b}^{l,i}_{(i,j+1)}.$$

if $r < s$, since $m_b = m_{b''}$, then the action of $E_{\mathcal{A}}$ for b'' is given by (a2). Then we have

$$E_{\mathcal{A}}b'' = \left(\tilde{f}_1^{\mathcal{A}}\right)^{p+s} \left(\tilde{f}_0^{\mathcal{A}}\right)^{p+1} \overline{b}^{l,i}_{(i,j+1)}.$$

Therefore we have,

$$\tilde{e}_1^{\mathcal{A}} E_{\mathcal{A}} \left(\left(\tilde{f}_0^{\mathcal{A}}\right)^{r} b \right) = E_{\mathcal{A}} \left(\left(\tilde{f}_0^{\mathcal{A}}\right)^{r} b'' \right).$$

If $r \geq s$, $m_b = 0, 1, 2$, since $m_{b'} < 0$, then the action of $E_{\mathcal{A}}$ for b' is given by (a3). Then we have

$$E_{\mathcal{A}}b' = \left(\tilde{f}_1^{\mathcal{A}}\right)^{p+s} \left(\tilde{f}_0^{\mathcal{A}}\right)^{p-1} \overline{b}^{l,i+1}_{(i+1,j-1)}.$$

therefore we have

$$\tilde{e}_1^{\mathcal{A}} E_{\mathcal{A}} \left(\left(\tilde{f}_0^{\mathcal{A}}\right)^{r} b \right) = E_{\mathcal{A}}^{s+l-2(i-s+1)} \left(\left(\tilde{f}_0^{\mathcal{A}}\right)^{r+1} \left(\tilde{f}_1^{\mathcal{A}}\right)^{p} \left(\tilde{f}_0^{\mathcal{A}}\right)^{p} \overline{b}^{l,k-s+1}_{(l-(i-s+1),j+s+1)} \right). \tag{5.3.15}$$

If $r \leq s$, $m_b \geq 3$, than the actions of $E_{\mathcal{A}}$ for b' is given by (a2), since $m_{b'} \geq 0$. Then we have,

$$E_{\mathcal{A}}b' = \left(\tilde{f}_1^{\mathcal{A}}\right)^{p+s} \left(\tilde{f}_0^{\mathcal{A}}\right)^{p} \overline{b}^{l,i}_{(i,j+1)},$$

(Figure 7). Thus we have

$$\begin{aligned} \tilde{e}_1^{\mathcal{A}} E_{\mathcal{A}} \left(\left(\tilde{f}_0^{\mathcal{A}}\right)^{r} b \right) &= E_{\mathcal{A}} \left(\left(\tilde{f}_0^{\mathcal{A}}\right)^{r+1} b' \right) \\ &= E_{\mathcal{A}}^{s+l-2(i-s+1)+m-2} \left(\left(\tilde{f}_0^{\mathcal{A}}\right)^{r+1} \left(\tilde{f}_1^{\mathcal{A}}\right)^{p-m+2} \left(\tilde{f}_0^{\mathcal{A}}\right)^{p-m+2} \overline{b}^{l,i-s+1}_{(l-(i-s+1),j-m+2+s)} \right) \end{aligned} \tag{5.3.16}$$

Therefore we have the case of (5.3.3), (5.3.4), (5.3.5), if $l-2i \leq l-i+2j-3q$.

The case of (a3)**:**$E_{\mathcal{A}}b \in B^{i+1}_{(i+1,j-1)}$**.**

This case is $b = E_{\mathcal{A}}^{n'} b_1$ $(l-2i < n' < l-i+2j-3q)$, $b = E_{\mathcal{A}}^{n'} b_2$ $(l-2i < n' < \left[\frac{j-i}{2}\right])$, $b = E_{\mathcal{A}}^{n'} b_3$ $(l-2i < n' < \left[\frac{j-i}{2}\right])$. (5.3.8).
Since $m_{b'} < m_{b''} = m_b < 0$, the action of $E_{\mathcal{A}}$ for b', b'' is given by (a3). Then we have

$$\begin{aligned} E_{\mathcal{A}}b &= \left(\tilde{f}_1^{\mathcal{A}}\right)^{p+s+1} \left(\tilde{f}_0^{\mathcal{A}}\right)^{p} \overline{b}^{l,i+1}_{(k+1,j-1)}, \\ E_{\mathcal{A}}b' &= \left(\tilde{f}_1^{\mathcal{A}}\right)^{p+s} \left(\tilde{f}_0^{\mathcal{A}}\right)^{p-1} \overline{b}^{l,i+1}_{(k+1,j-1)}, \\ E_{\mathcal{A}}b'' &= \left(\tilde{f}_1^{\mathcal{A}}\right)^{p+s} \left(\tilde{f}_0^{\mathcal{A}}\right)^{p} \overline{b}^{l,i+1}_{(k+1,j-1)}, \end{aligned}$$

(Figure 8). With Proposition 5.4, we have

$$\begin{aligned} \tilde{e}_1^{\mathcal{A}} E_{\mathcal{A}} \left(\left(\tilde{f}_0^{\mathcal{A}}\right)^{r} b \right) &= E_{\mathcal{A}} \left(\left(\tilde{f}_0^{\mathcal{A}}\right)^{r} b'' \right) \qquad (r < s+1), \\ \tilde{e}_1^{\mathcal{A}} E_{\mathcal{A}} \left(\left(\tilde{f}_0^{\mathcal{A}}\right)^{r} b \right) &= E_{\mathcal{A}} \left(\left(\tilde{f}_0^{\mathcal{A}}\right)^{r+1} b' \right) \quad (r \geq s+1). \end{aligned} \tag{5.3.17}$$


$\overline{b}^{l,i}_{(i,j)}$
p–1
p–1
s
p
b'
b''
b
s–1
$\swarrow \tilde{f}_1^{\mathcal{A}}$
$\searrow \tilde{f}_0^{\mathcal{A}}$

Figure 6: Part II, the location of b, b' and b'' in the crystal graph $B^i_{(i,j)}$.


$\overline{b}^{l,i}_{(i,j+1)}$
p
p
s
p+1
$E_{\mathcal{A}}b''$
$E_{\mathcal{A}}b$
s–1
$\swarrow \tilde{f}_1^{\mathcal{A}}$
$\searrow \tilde{f}_0^{\mathcal{A}}$

Figure 7: Part II, the case of (a2), the location of $E_{\mathcal{A}}b$ and $E_{\mathcal{A}}b''$ in the crystal graph of $B^i_{(i,j+1)}$.

Therefore we have the case of $(l-2i<n\le l-i+2j-3q)$ of (5.3.2), the case of $(l-2i<n\le \left[\frac{j-i}{2}\right])$ of (5.3.6) and (5.3.7),

By (5.3.9) – (5.3.17), we have Proposition 5.30. □

5.3.2 Actions of $\tilde{e}_1^{\mathcal{G}}$

Proposition 5.31 For $b\in\mathcal{G}$, $n\in\mathbf{Z}_{>0}$ if $\tilde{f}_1^{\mathcal{G}}\left(\tilde{e}_2^{\mathcal{G}}\right)^n b=\left(\tilde{e}_2^{\mathcal{G}}\right)^n \tilde{f}_1^{\mathcal{G}} b\neq 0$, we have

$$\tilde{f}_1^{\mathcal{G}}\left(\tilde{e}_2^{\mathcal{G}}\right)^k b=\left(\tilde{e}_2^{\mathcal{G}}\right)^k \tilde{f}_1^{\mathcal{G}} b\ \ (0\le k\le n).$$

Proof We prove the proposition by the induction on k. We assume

$$\tilde{f}_1\left(\tilde{e}_2\right)^{k'} b=\left(\tilde{e}_2\right)^{k'}\tilde{f}_1 b\ \ (0\le k'<k),$$

then we prove

$$\tilde{f}_1\tilde{e}_2\left(\tilde{e}_2\right)^{k-1} b=\tilde{e}_2\tilde{f}_1\left(\tilde{e}_2\right)^{k-1} b. \tag{5.3.18}$$

We write $\tilde{e}_2^{k-1}b=b_1\otimes\cdots\otimes b_l$. We have

$$\tilde{f}_1\left(\tilde{e}_2\right)^{k-1} b=b_1\otimes\cdots\otimes\tilde{f}_1 b_{k_1}\otimes\cdots\otimes b_l, \tag{5.3.19}$$

$$\tilde{e}_2\left(\tilde{e}_2\right)^{k-1} b=b_1\otimes\cdots\otimes\tilde{e}_1 b_{k_2}\otimes\cdots\otimes b_l, \tag{5.3.20}$$

where $1\le k_1,k_2\le l$.

The case of $k_1>k_2$. Similar to Proposition 5.6, by Proposition 3.3 we see that the action of f_1 increase the number of u_+ of $\mathrm{Red}_2(b)$. Then the operator $\tilde{f}_1$ does not influence the action of $\tilde{e}_2$. In a similar way, we see that the operator $\tilde{e}_2$ does not influence the action of $\tilde{f}_1$. Therfore we have (5.3.18).



$\overline{b}^{l,i+1}_{(i,j+1)}$
p–1
p–1
s+1
p
$E_{\mathcal{A}}b'$
$E_{\mathcal{A}}b''$
$E_{\mathcal{A}}b$
s
$\swarrow \tilde{f}_1^{\mathcal{A}}$ $\searrow \tilde{f}_0^{\mathcal{A}}$

Figure 8: Part II, the case of (a3), the location of $E_{\mathcal{A}}b$, $E_{\mathcal{A}}b'$ and $E_{\mathcal{A}}b''$ in the crystal graph of $B^{i+1}_{(i,j+1)}$.

The case of $k_1 \leq k_2$. We assume

$$\tilde{f}_1\tilde{e}_2\left(\tilde{e}_2\right)^{k-1}b \neq \tilde{e}_2\tilde{f}_1\left(\tilde{e}_2\right)^{k-1}b. \tag{5.3.21}$$

By assumption, at most one of following relations are satisfied:

$$\tilde{f}_1\tilde{e}_2\left(\tilde{e}_2^{k-1}\right)b = \begin{cases} b_1\otimes\cdots\otimes\tilde{f}_1\tilde{e}_2b_{k_2}\otimes\cdots\otimes b_l \\ b_1\otimes\cdots\otimes\tilde{e}_2b_{k_2}\otimes\cdots\otimes\tilde{f}_1b_{k_1'}\otimes\cdots\otimes b_l \end{cases}, \tag{5.3.22}$$

$$\tilde{e}_2\tilde{f}_1\left(\tilde{e}_2^{k-1}\right)b = \begin{cases} b_1\otimes\cdots\otimes\tilde{e}_2\tilde{f}_1b_{k_1}\otimes\cdots\otimes b_l \\ b_1\otimes\cdots\otimes\tilde{e}_2b_{k_2'}\otimes\cdots\otimes\tilde{f}_1b_{k_1}\otimes\cdots\otimes b_l \end{cases}, \tag{5.3.23}$$

Since $\tilde{f}_1u_+ = u_-$ and $\Psi_1(b) = u_-^{\varepsilon_1(b)}u_+^{\varphi_1(b)}$, we have $k_1 < k_1'$. In a similar way we have $k_2' < k_2$. If (5.3.22) is satisfied, we see that k_1-th element of $\tilde{e}_2^{k'}\tilde{f}_1\tilde{e}_2^{k}b$ and $\tilde{f}_1\tilde{e}_2^{k'}\tilde{e}_2^{k}b$ $(1 \leq k' \leq n-k)$ are different. Then we have

$$\tilde{e}_2^{n}\tilde{f}_1b \neq \tilde{f}_1\tilde{e}_2^{n}b. \tag{5.3.24}$$

This is contradiction. In a similar way, if (5.3.23) is satisfied, we have contradiction. Thus we have (5.3.18). □

We prove from (5.3.2) to (5.3.8) exchanging $\tilde{e}_1^{\mathcal{A}}$ with $\tilde{e}_1^{\mathcal{G}}$. For example, we denote following equations exchanging $\tilde{e}_1^{\mathcal{A}}$ with $\tilde{e}_1^{\mathcal{G}}$ from (5.3.2) to (5.3.5). We put $b_1 = \left(\tilde{f}_1^{\mathcal{A}}\right)^q\left(\tilde{f}_0^{\mathcal{A}}\right)^p\overline{b}^{l,i}_{(l-i,j)} \in B_C$ $(3q \geq i+2j-\left[\frac{j-i}{2}\right]$, $p+2q > i+2j)$.

We are going to show

$$\begin{aligned} \tilde{e}_1^{\mathcal{G}}\left(\Phi\left(E_{\mathcal{A}}^{n}\,b_1\right)\right) &= \Phi\left(E_{\mathcal{A}}^{n}\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q-1}\left(\tilde{f}_0^{\mathcal{A}}\right)^{p}\overline{b}^{l,i}_{(l-i,j)}\right)\right), && (5.3.25)\\ \tilde{e}_1^{\mathcal{G}}\left(\Phi\left(E_{\mathcal{A}}^{l-i-j+3(j-q)+1}\,b_1\right)\right) &= \Phi\left(E_{\mathcal{A}}^{l-i-j+3(j-q)}\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q}\left(\tilde{f}_0^{\mathcal{A}}\right)^{p+1}\overline{b}^{l,i}_{(l-i,j+1)}\right)\right), && (5.3.26)\end{aligned}$$

$$\tilde{e}_1^{\mathcal{G}}\left(\Phi\left(E_{\mathcal{A}}^{l-i-j+3(j-q)+2}\ b_1\right)\right) = \Phi\left(E_{\mathcal{A}}^{l-i-j+3(j-q)}\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q+1}\left(\tilde{f}_0^{\mathcal{A}}\right)^{p+2}\overline{b}^{l,i}_{(l-i,j+2)}\right)\right), \tag{5.3.27}$$

$$\tilde{e}_1^{\mathcal{G}}\left(\Phi\left(E_{\mathcal{A}}^{l-i-j+3(j-q+1)+n'}\ b_1\right)\right) = \Phi\left(E_{\mathcal{A}}^{l-i-j+3(j-q)+n'}\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q+2}\left(\tilde{f}_0^{\mathcal{A}}\right)^{p+3}\overline{b}^{l,i}_{(l-i,j+3)}\right)\right), \tag{5.3.28}$$

where $0 < n < l-j-i+3(j-q)$, $0 < n' < 3(y-1)$.

Here we prove the case of $l-i-j \equiv 0 \pmod 3$, we can prove other cases similarly.

By Proposition 5.20, we have

$$\Phi(b_1) = \left\llbracket \llbracket 1 \rrbracket^i \llbracket 6 \rrbracket^{j-q} \llbracket 6 \rrbracket^y \llbracket \mathcal{C} \rrbracket^{y+q+p-j-i} \llbracket 2 \rrbracket^{y-1} \llbracket \overline{1} \rrbracket^{j+i-p} \right\rrbracket.$$

The case of (5.3.25), $n=0$. By (E1), we have

$$\tilde{e}_1^{\mathcal{G}}\left(\Phi(b_1)\right) = \Phi\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q-1}\left(\tilde{f}_0^{\mathcal{A}}\right)^{p}\overline{b}^{l,i}_{(l-i,j)}\right).$$

The case of (5.3.25), $n = l-i-j+3(j-q)$. Put $b_4 = E_{\mathcal{A}}^{l-j-i+3(j-q)} b$. We see

$$\Phi(b_4) = \left\llbracket \llbracket 1 \rrbracket^i \llbracket 2 \rrbracket^{j-p} \llbracket 2 \rrbracket^y \llbracket \mathcal{C} \rrbracket^{y+q+p-j-i} \llbracket \overline{2} \rrbracket^{y} \llbracket \overline{1} \rrbracket^{j+i-p} \right\rrbracket.$$

Using Definition 3.4

$$\begin{array}{rcl} \Psi_1(b_4) & = & u_-^{j-p+i} u_+^y u_-^{y+q+p-j-i} u_+^{y+q+p-j-i} u_-^{y+j-p} u_+^i, \\ \mathrm{Red}_1(b_4) & = & u_-^{j-p+i} u_-^{q+p-j-i} u_+^{2q+p-2j-i} u_+^i. \end{array}$$

By Proposition 3.9, we have

$$\begin{array}{rcl} \tilde{e}_1^{\mathcal{G}}\left(\Phi(b_4)\right) & = & \left\llbracket \llbracket 1 \rrbracket^i \llbracket 2 \rrbracket^{j-p} \llbracket 2 \rrbracket^y \, \tilde{e}_1^{\mathcal{G}} \llbracket \mathcal{C} \rrbracket^{y+q+p-j-i} \llbracket \overline{2} \rrbracket^{y} \llbracket \overline{1} \rrbracket^{j+i-p} \right\rrbracket \\ & = & \left\llbracket \llbracket 1 \rrbracket^i \llbracket 2 \rrbracket^{j-p} \llbracket 2 \rrbracket^y \llbracket 6 \rrbracket \llbracket \mathcal{C} \rrbracket^{y+q+p-j-i-1} \llbracket \overline{2} \rrbracket^{y} \llbracket \overline{1} \rrbracket^{j+i-p} \right\rrbracket. \end{array}$$

Put $b_4' = E_{\mathcal{A}}^n\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q-1}\left(\tilde{f}_0^{\mathcal{A}}\right)^{p}\overline{b}^{l,i}_{(l-i,j)}\right)$. We see

$$\Phi(b_4') = \left\llbracket \llbracket 1 \rrbracket^i \llbracket 2 \rrbracket^{j-p} \llbracket 2 \rrbracket^y \llbracket 6 \rrbracket \llbracket \mathcal{C} \rrbracket^{y+p+q-j-i-1} \llbracket \overline{2} \rrbracket^{y} \llbracket \overline{1} \rrbracket^{j+i-p} \right\rrbracket.$$

Thus we have

$$\tilde{e}_1^{\mathcal{G}}\left(\Phi\left(E_{\mathcal{A}}^n b_1\right)\right) = \Phi\left(E_{\mathcal{A}}^n\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q-1}\left(\tilde{f}_0^{\mathcal{A}}\right)^{p}\overline{b}^{l,i}_{(l-i,j)}\right)\right).$$

The case of (5.3.26). We set $b_5 = E_{\mathcal{A}}^{l-j-i+3(j-q)+1} b$.

$$\Phi(b_5) = \left\llbracket \llbracket 1 \rrbracket^i \llbracket 2 \rrbracket^{j-p} \llbracket 2 \rrbracket^y \llbracket \mathcal{C} \rrbracket^{y+q+p-j-i} \llbracket \overline{3} \rrbracket \llbracket \overline{2} \rrbracket^{y-1} \llbracket \overline{1} \rrbracket^{j+i-p} \right\rrbracket.$$

We consider $\tilde{e}_1^{\mathcal{G}}(\Phi(b_5))$ with u_+ and u_-.

$$\begin{array}{rcl} \Psi_1(\Phi(b_5)) & = & u_-^{j+i-p} u_+^{y-1} u_0 u_-^{y+q+p-j-i} u_+^{y+q+p-j-i} u_-^{y+j-p} u_+^i, \\ \mathrm{Red}_1\left(\Phi(b_5)\right) & = & u_-^{j+i-p} u_-^{p+q-j-i+1} u_+^{2q+p-2j-i} u_+^i. \end{array}$$

Then we have

$$\begin{aligned}\tilde{e}_1^{\mathcal{G}}\left(\Phi(b_5)\right) &= \Big\llbracket \llbracket 1 \rrbracket^{i} \llbracket 2 \rrbracket^{j-p} \llbracket 2 \rrbracket^{y}\, \tilde{e}_1^{\mathcal{G}} \llbracket \mathcal{C} \rrbracket^{y+q+p-j-i} \llbracket \overline{3} \rrbracket \llbracket \overline{2} \rrbracket^{y-1} \llbracket \overline{1} \rrbracket^{j+i-p} \Big\rrbracket \\ &= \Big\llbracket \llbracket 1 \rrbracket^{i} \llbracket 2 \rrbracket^{j-p} \llbracket 2 \rrbracket^{y} \llbracket 6 \rrbracket \llbracket \mathcal{C} \rrbracket^{y+q+p-j-i-1} \llbracket \overline{3} \rrbracket \llbracket \overline{2} \rrbracket^{y-1} \llbracket \overline{1} \rrbracket^{j+i-p} \Big\rrbracket .\end{aligned}$$

We put $b_5' = E_{\mathcal{A}}^{l-i-j+3(j-q)} \left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q} \left(\tilde{f}_0^{\mathcal{A}}\right)^{p+1} \overline{b}^{\,l,i}_{(l-i,j+1)} \right)$. Then we have

$$\Phi(b_5') = \Big\llbracket \llbracket 1 \rrbracket^{i} \llbracket 2 \rrbracket^{j-q} \llbracket 2 \rrbracket^{y} \llbracket 6 \rrbracket \llbracket \mathcal{C} \rrbracket^{y+p+q-j-i-1} \llbracket \overline{3} \rrbracket \llbracket \overline{2} \rrbracket^{y-1} \llbracket \overline{1} \rrbracket^{j+i-p} \Big\rrbracket .$$

Thus we have

$$\tilde{e}_1^{\mathcal{G}} \left(\Phi \left(E_{\mathcal{A}}^{l-i-j+3(j-q)+1} b_1 \right) \right) = E_{\mathcal{A}}^{l-i-j+3(j-q)} \left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q} \left(\tilde{f}_0^{\mathcal{A}}\right)^{p+1} \overline{b}^{\,l,i}_{(l-i,j+1)} \right).$$

Similarly, in case of (5.3.27), $n' = 0$ and $n' = 2(l-i-j) + 3(j-q)$ of (5.3.28), we have

$$\tilde{e}_1^{\mathcal{G}} \left(\Phi \left(E_{\mathcal{A}}^{l-i-j+3(j-q)+2} b_1 \right) \right) = \Phi \left(E_{\mathcal{A}}^{l-i-j+3(j-q)} \left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q+1} \left(\tilde{f}_0^{\mathcal{A}}\right)^{p+2} \overline{b}^{\,l,i}_{(l-i,j+2)} \right) \right),$$

$$\tilde{e}_1^{\mathcal{G}} \left(\Phi \left(E_{\mathcal{A}}^{l-i-j+3(j-p)+3} b_1 \right) \right) = \Phi \left(E_{\mathcal{A}}^{l-i-j+3(j-q)} \left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q+2} \left(\tilde{f}_0^{\mathcal{A}}\right)^{p+3} \overline{b}^{\,l,i}_{(l-i,j+3)} \right) \right),$$

$$\tilde{e}_1^{\mathcal{G}} \left(\Phi \left(E_{\mathcal{A}}^{2(l-i-j)+3(j-p)} b_1 \right) \right) = \Phi \left(E_{\mathcal{A}}^{2(l-i-j)+3(j-q-1)} \left(\left(\tilde{f}_1^{\mathcal{A}}\right)^{q+2} \left(\tilde{f}_0^{\mathcal{A}}\right)^{p+3} \overline{b}^{\,l,i}_{(l-i,j+3)} \right) \right).$$

By Proposition 5.31, we have $\tilde{e}_1^{\mathcal{G}}\Phi = \Phi\tilde{e}_1^{\mathcal{A}}$ for $E_{\mathcal{A}}^{n} b_1$ $(0 \le n \le \varepsilon_{\mathcal{A}}(b_1) = 2(l-i-j) + 3(j-p))$.

Similarly, we can prove $\tilde{e}_1^{\mathcal{G}}\Phi = \Phi\tilde{e}_1^{\mathcal{A}}$ for any $b \in \mathcal{A}$.

5.4 Selection of minimal

5.4.1 Selection of minimal elements in $B^{G_2}(l\Lambda_1)$

In this section, we only consider crystal B^l, so we denote $\tilde{e}_1$, $\tilde{e}_2$ instead of $\tilde{e}_1^{\mathcal{G}}$, $\tilde{e}_2^{\mathcal{G}}$ respectively for simplicity. By definition 3.16, minimal elements are $b \in B^l$ such that $\langle c, \varphi(b) \rangle = l$. At first we search for the element $b \in B(l\Lambda)$ such that $\langle c, \varphi_1(b)\Lambda_1 + \varphi_1(b)\Lambda_2 \rangle = l$. Next we verify that there does not exist $b \in B(l\Lambda_1)$ such that $\langle c, \varphi_1(b)\Lambda_1 + \varphi_2(b)\Lambda_2 \rangle < l$. Then we verify that for $b \in B^l$ such that $\langle c, \varphi_1(b)\Lambda_1 + \varphi_2(b)\Lambda_2 \rangle = l$, we have $\varphi_0(b) = 0$

Proposition 5.32 *For $b \in B^{G_2}(l\Lambda_1)$ such that $\varepsilon_2(b) = 0$, we have*

$$\min\{2\varphi_1(\tilde{f}_2^{(k)}(b)) + \varphi_2(\tilde{f}_2^{(k)}(b)) \mid k = 0, \ldots, \varphi_2(b)\} \ge l - \left[\frac{1}{2} wt_0(b)\right],$$

Proof We consider

$$\min\{2\varphi_1(b') + \varphi_2(b') \mid b' = \tilde{f}_2^{(k)}(b),\ k = 0, \ldots, \varphi_2(b)\}. \tag{5.4.1}$$

By Proposition 5.6, we have

$$2\varphi_1(\tilde{f}_2^{(k+1)}(b)) + \varphi_2(\tilde{f}_2^{(k+1)}(b)) - (2\varphi_1(\tilde{f}_2^{(k)}(b)) + \varphi_2(\tilde{f}_2^{(k)}(b))) = \left\{ \begin{array}{ll} 1 & (k < k') \\ -1 & (k \le k') \end{array} \right. .$$

We set $b' = \tilde{f}_2^{\varphi_2(b)} b$. Then (5.4.1) is

$$2\varphi_1(b) + \varphi_2(b) - \frac{\varphi_2(b) - (2\varphi_1(b') - (2\varphi_1(b) + \varphi_2(b)))}{2} = \varphi_1(b) + \varphi_1(b').$$

Then we prove

$$\varphi_1(b) + \varphi_1(b') - l + \left[\frac{1}{2}\mathrm{wt}_0(b)\right] \geq 0.$$

We recall that $x_i(b) = \sharp\left\{ \boxed{b_k} = \boxed{i} \;\middle|\; k = 1, \ldots, n\right\}$ ($i = 1, \ldots, 6, 0_1, 0_2$), $\overline{x}_i(b) = \sharp\left\{ \boxed{b_k} = \boxed{\overline{i}} \;\middle|\; k = 1, \ldots, n\right\}$ ($i = 1, \ldots, 6$). We consider b such that $\varepsilon_2(b) = 0$. Such an element b satisfy following conditions:

(1) $0 \leq x_1(b) + \overline{x}_1(b) \leq l$. This is because $\tilde{f}_2\left(\boxed{1}\right) = 0$, $\tilde{e}_2\left(\boxed{1}\right) = 0$, $\tilde{f}_2\left(\boxed{\overline{1}}\right) = 0$, $\tilde{e}_2\left(\boxed{\overline{1}}\right) = 0$.

(2) $x_{0_2}(b) = 0$ or 1. Since $\tilde{f}_2\left(\boxed{0_2}\right) = 0$, $\tilde{e}_2\left(\boxed{0_2}\right) = 0$ and by Proposition 3.15 $x_{0_2}(b) < 2$.

(3) $\overline{x}_2(b) = \overline{x}_3(b) = \overline{x}_4(b) = \overline{x}_5(b) = \overline{x}_{0_1}(b) = 0$. Because if $\overline{x}_2(b) > 0$ or $\overline{x}_3(b) > 0$ or $\overline{x}_4(b) > 0$ or $\overline{x}_5(b) > 0$ or $\overline{x}_{0_1}(b) > 0$ then $\varepsilon_2(b) > 0$.

(4) $x_3(b) + x_4(b) + x_5(b) = 0$ or 1. By Proposition 3.15.

(5) $x_6(b) \leq \overline{x}_6(b)$. In particular, if $x_3(b) = 1$ or $x_4(b) = 1$ then $x_6(b) < \overline{x}_6(b)$ and if $x_5(b)$ then $x_6(b) = 0$.

(6) $\displaystyle\sum_{i=1}^{6} (x_i(b) + \overline{x}_i(b)) + x_{0_1}(b) + x_{0_2}(b) = l$.

We start with the case of $x_5(b) = 0$.
By calculation, we have $x_1(b') = x_1(b)$, $x_6(b') = x_2(b) + x_3(b) + x_4(b) + x_6(b)$, $x_{0_2}(b') = x_{0_2}(b)$, $\overline{x}_6(b') = x_6(b)$, $\overline{x}_4(b') = x_4(b)$, $\overline{x}_3(b') = x_3(b)$, $\overline{x}_2(b') = \overline{x}_6(b) - x_{0_2}(b) - x_6(b) - x_4(b) - x_3(b)$, $\overline{x}_1(b') = \overline{x}_1(b)$. We put $\delta = \mathrm{wt}_0(b) \bmod 2$. Thus we have

$$\begin{aligned}
-wt_0(b) &= 2x_1(b) + x_2(b) + x_3(b) + x_4(b) + x_6(b) - \overline{x}_6(b) - 2\overline{x}_1(b), \\
l &= x_1(b) + x_2(b) + x_3(b) + x_4(b) + x_6(b) + x_{0_2}(b) + \overline{x}_6(b) + \overline{x}_1(b), \\
l - \left[\frac{1}{2}wt_0(b)\right] &= 2x_1(b) + \frac{3}{2}x_2(b) + \frac{1}{2}x_3(b) + \frac{1}{2}x_4(b) + \frac{3}{2}x_6(b) + \frac{1}{2}\overline{x}_6(b) + \frac{\delta}{2}, \\
\varphi_1(b) &= x_1(b) + (x_{0_2}(b) + 2x_6(b) + x_4(b) - x_2(b))_+, \\
\varphi_1(b') &= x_1(b') + x_{0_2}(b') + 2x_6(b') + (\overline{x}_2(b') - \overline{x}_4(b') - \overline{x}_6(b') - x_{0_2}(b'))_+, \\
&= x_1(b) + 2x_2(b) + 2x_3(b) + 2x_4(b) + 2x_6(b) + x_{0_2}(b) \\
&\quad + (\overline{x}_6(b) - x_{0_2}(b) - 3x_6(b) - 2x_4(b) - x_3(b))_+.
\end{aligned}$$

$$\begin{aligned}
\varphi_1(b) + \varphi_1(b') - l + \left[\frac{1}{2}wt_0(b)\right] &= \frac{1}{2}\left|x_{0_2}(b) + 2x_6(b) + x_4(b) - x_2(b)\right| \\
&\quad + \frac{1}{2}\left|\overline{x}_6(b) - x_{0_2}(b) - 3x_6(b) - 2x_4(b) - x_3(b)\right| - \frac{\delta}{2} \qquad (5.4.2) \\
&\geq 0 \text{ (integer greater than } -\tfrac{1}{2} \text{)}.
\end{aligned}$$

The case of $x_5(b) = 1$. By Proposition 3.15, $x_3(b) = x_4(b) = x_6(b) = x_{0_2}(b) = 0$. Similar to the case of $x_5(b) = 0$, we have

$$\varphi_1(b) + \varphi_1(b') - l + \left[\frac{1}{2}wt_0(b)\right] = \frac{1}{2}x_2(b) + \frac{1}{2}\overline{x}_6(b) - \frac{\delta}{2} \geq 0 \text{ (integer greater than } -\tfrac{1}{2} \text{)} \qquad (5.4.3)$$

□

A minimal element is an element $b \in B^l$ such that $\langle c, \varphi(b)\rangle = l$. By $\langle c, \varphi(b)\rangle = \varphi_0(b) + 2\varphi_1(b) + \varphi_2(b)$ and Proposition 5.32, if $\mathrm{wt}_0(b) > 0$ we have $\langle c, \varphi(b)\rangle > l$. Therefore for a minimal element b we must have $\mathrm{wt}_0(b) = 0$.

Lemma 5.33 *we have*

$$\min\{\langle c, \varphi(b)\rangle \mid b \in B(n\Lambda_1)\} = l.$$

Proof By (5.4.2) and (5.4.3), the elements b such that $\varepsilon_2(b) = 0$, $\varphi_1(b) + \varphi_1(\tilde{f}_2^{\varphi_2(b)}(b)) = l$ satisfy following conditions.
If $x_5(b) = 0$,

$$\begin{aligned} x_2(b) &= 2x_6(b) + x_{0_2}(b) + x_4(b), \\ \overline{x}_2(b) &= 3x_6(b) + x_{0_2}(b) + 2x_4(b) + x_3(b). \end{aligned}$$

If $x_5(b) = 1$,

$$\begin{aligned} x_2(b) &= 0, \\ x_6(b) &= 0. \end{aligned}$$

Then we can express b such that $\varepsilon_2(b) = 0$, $\varphi_1(b) + \varphi_1(\tilde{f}_2^{\varphi_2(b)}(b)) = l$ as:

$$b = \begin{cases} \Big\llbracket \llbracket 1 \rrbracket^m \llbracket 2 \rrbracket^n \llbracket \mathcal{C} \rrbracket^n \llbracket \overline{6} \rrbracket^n \llbracket \overline{1} \rrbracket^m \Big\rrbracket & (l = 2m + 3n), \\ \Big\llbracket \llbracket 1 \rrbracket^m \llbracket 2 \rrbracket^n \llbracket 4 \rrbracket \llbracket \mathcal{C} \rrbracket^{n-1} \llbracket \overline{6} \rrbracket^{n+1} \llbracket \overline{1} \rrbracket^m \Big\rrbracket & (n > 0,\ l = 2m + 3n + 1), \\ \Big\llbracket \llbracket 1 \rrbracket^m \llbracket 5 \rrbracket \llbracket \overline{1} \rrbracket^m \Big\rrbracket & (l = 2m + 1), \\ \Big\llbracket \llbracket 1 \rrbracket^m \llbracket 2 \rrbracket^n \llbracket 3 \rrbracket \llbracket \mathcal{C} \rrbracket^n \llbracket \overline{6} \rrbracket^{n+1} \llbracket \overline{1} \rrbracket^m \Big\rrbracket & (l = 2m + 3n + 2). \end{cases}$$

where $m, n \in \mathbf{Z}_{\geq 0}$. We put

$$\tilde{b}_{(m)} = \tilde{f}_2^{3n} b = \begin{cases} \Big\llbracket \llbracket 1 \rrbracket^m \llbracket 2 \rrbracket^n \llbracket \mathcal{C} \rrbracket^n \llbracket \overline{2} \rrbracket^n \llbracket \overline{1} \rrbracket^m \Big\rrbracket & (l = 2m + 3n), \\ \Big\llbracket \llbracket 1 \rrbracket^m \llbracket 2 \rrbracket^n \llbracket 4 \rrbracket \llbracket \mathcal{C} \rrbracket^{n-1} \llbracket \overline{4} \rrbracket \llbracket \overline{6} \rrbracket^n \llbracket \overline{1} \rrbracket^m \Big\rrbracket & (n > 0, l = 2m + 3n + 1), \\ \Big\llbracket \llbracket 1 \rrbracket^m \llbracket 0_1 \rrbracket \llbracket \overline{1} \rrbracket^m \Big\rrbracket & (l = 2m + 1), \\ \Big\llbracket \llbracket 1 \rrbracket^m \llbracket 2 \rrbracket^n \llbracket 3 \rrbracket \llbracket \mathcal{C} \rrbracket^n \llbracket \overline{3} \rrbracket \llbracket \overline{2} \rrbracket^n \llbracket \overline{1} \rrbracket^m \Big\rrbracket & (l = 2m + 3n + 2), \end{cases}$$

where $m \in \mathbf{Z}_{\geq} 0$ such taht $l - 2m \geq 0$. We prove $\langle c, 2\varphi_1(\tilde{b}_{(m)})\Lambda_1 + \varphi_2(\tilde{b}_{(m)})\Lambda_2\rangle = l$ $(k = 1, 2, 3)$. We prove the case of $l = 2m + 3n$. By Proposition 3.9

$$\begin{aligned} \Psi_1(\tilde{b}_{(m)}) &= u_-^m u_+^n u_-^n u_+^n u_-^n u_+^m, \\ \mathrm{Red}_1\left(\tilde{b}_{(m)}\right) &= u_-^m u_+^m, \end{aligned}$$

we have

$$\varphi_1(\tilde{b}_{(m)}) = m.$$

By Proposition 3.9

$$\begin{aligned} \Psi_2(\tilde{b}_{(m)}) &= u_0 u_-^{3n} u_0 u_+^{2n} u_0, \\ \mathrm{Red}_2\left(\tilde{b}_{(m)}\right) &= u_-^{3n} u_+^{3n}, \end{aligned}$$

we have

$$\varphi_2(\tilde{b}_{(m)}) = 3n.$$

Therefore,

$$\langle c, 2\varphi_1(\tilde{b}_{(m)})\Lambda_1 + \varphi_2(\tilde{b}_{(m)})\Lambda_2\rangle = 2m + 3n.$$

In similar way, we can prove other cases.

Let $B_l^0 \cong \{b \mid b \in B(l\Lambda_1),\ \langle c, \varphi_1(b)\Lambda_1 + \varphi_2(b)\rangle = l\}$, $(P_{cl}^+)_l^0 \cong \{\lambda \in \mathbf{Z}\Lambda_1 + \mathbf{Z}\Lambda_2 \mid \langle c, \lambda\rangle = l\}$. Then we see following proposition easily.

Proposition 5.34 *The maps $\varepsilon,\ \varphi\ :\ B_l^0 \to (P_{cl}^+)_l^0$ are bijective.*

5.4.2 Existence of minimal elements on $B^{G_2}(l\Lambda_1)$

We consider $\left(\tilde{f}_1^{\mathcal{A}}\right)^i \left(\tilde{f}_0^{\mathcal{A}}\right)^i \overline{b}^{l,i}_{(i,i)}$. By Definition 5.12, we have

$$\left(\tilde{f}_1^{\mathcal{A}}\right)^i \left(\tilde{f}_0^{\mathcal{A}}\right)^i \overline{b}^{l,i}_{(i,i)} = \left(\tilde{e}_2^{\mathcal{A}}\right)^{l-2i} \left(\tilde{f}_1^{\mathcal{A}}\right)^i \left(\tilde{f}_0^{\mathcal{A}}\right)^i \overline{b}^{l,i}_{((l-i),i)}.$$

For $b \in B^i_{k\Lambda_1 + j\Lambda_0}$, we define $y = y(b) = \left[\frac{l-i-j}{3}\right]$.
By Proposition 5.20 and Lemma 5.21,

$$\Phi\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^i \left(\tilde{f}_0^{\mathcal{A}}\right)^i \overline{b}^{l,i}_{((l-i),i)}\right) =$$
$$\begin{cases} \left\llbracket \llbracket 1 \rrbracket^i \llbracket 6 \rrbracket^y \llbracket \mathcal{C} \rrbracket^y \llbracket \overline{2} \rrbracket^y \llbracket \overline{1} \rrbracket^i \right\rrbracket & (l-2i \equiv 0 \pmod 3), \\ \left\llbracket \llbracket 1 \rrbracket^i \llbracket \overline{5} \rrbracket \llbracket \overline{1} \rrbracket^i \right\rrbracket & (l-2i \equiv 1 \pmod 3), y = 0), \\ \left\llbracket \llbracket 1 \rrbracket^i \llbracket 6 \rrbracket^y \llbracket 6 \rrbracket \llbracket \mathcal{C} \rrbracket^{y-1} \llbracket \overline{4} \rrbracket \llbracket \overline{2} \rrbracket^y \llbracket \overline{1} \rrbracket^i \right\rrbracket & (l-2i \equiv 1 \pmod 3), y > 0), \\ \left\llbracket \llbracket 1 \rrbracket^i \llbracket 6 \rrbracket^y \llbracket 6 \rrbracket \llbracket \mathcal{C} \rrbracket^y \llbracket \overline{3} \rrbracket \llbracket \overline{2} \rrbracket^y \llbracket \overline{1} \rrbracket^i \right\rrbracket & (l-2i \equiv 2 \pmod 3)), \end{cases}$$

we have

$$\Phi\left(\left(\tilde{e}_2^{\mathcal{A}}\right)^{l-2i} \left(\tilde{f}_1^{\mathcal{A}}\right)^i \left(\tilde{f}_0^{\mathcal{A}}\right)^i \overline{b}^{l,i}_{((l-i),i)}\right) =$$
$$\begin{cases} \left\llbracket \llbracket 1 \rrbracket^i \llbracket 2 \rrbracket^y \llbracket \mathcal{C} \rrbracket^y \llbracket \overline{2} \rrbracket^y \llbracket \overline{1} \rrbracket^i \right\rrbracket & (l-2i \equiv 0 \pmod 3), \\ \left\llbracket \llbracket 1 \rrbracket^i \llbracket 0_1 \rrbracket \llbracket \overline{1} \rrbracket^i \right\rrbracket & (l-2i \equiv 1 \pmod 3), y = 0), \\ \left\llbracket \llbracket 1 \rrbracket^i \llbracket 2 \rrbracket^y \llbracket 4 \rrbracket \llbracket \mathcal{C} \rrbracket^{y-1} \llbracket \overline{4} \rrbracket \llbracket \overline{2} \rrbracket^y \llbracket \overline{1} \rrbracket^i \right\rrbracket & (l-2i \equiv 1 \pmod 3), y > 0), \\ \left\llbracket \llbracket 1 \rrbracket^i \llbracket 2 \rrbracket^y \llbracket 3 \rrbracket \llbracket \mathcal{C} \rrbracket^y \llbracket \overline{3} \rrbracket \llbracket \overline{2} \rrbracket^y \llbracket \overline{1} \rrbracket^i \right\rrbracket & (l-2i \equiv 2 \pmod 3)), \end{cases}$$

Therefore we have proved existence of minimal elements. By (5.1.12) we see

$$\varphi_0\left(\left(\tilde{f}_1^{\mathcal{A}}\right)^i \left(\tilde{f}_0^{\mathcal{A}}\right)^i \overline{b}^{l,i}_{(i,i)}\right) = 0.$$

Thus we have

$$\langle c, \varphi(\tilde{b}_{(m)})\rangle = l,$$

where $m \in \mathbf{Z}_{\geq 0}$.

5.4.3 Selection of minimals on B^l

By (5.2.6), for $n < l$, we have

$$\varphi_0\Big|_{B^{n+1}}(b) = \varphi_0\Big|_{B^n}(b) + 1.$$

Then for $b \in B^n$ such that $\langle c, \varphi(b)\rangle = n$, we have $\langle c, \varphi(b)\rangle = n + 1$ on B^{n+1}. If $l = 1$, by calculation we see $\min\left\{\langle c, \varphi(b)\rangle \mid b \in B^1\right\} = 1$. Then we see

$$\min\left\{\langle c, \varphi(b)\rangle \;\middle|\; b \in B^l\right\} = l,$$

inductively. Therefore a minimal element for level $l - 1$ is also a minimal element for level l. Thus it is obvious that ε and φ are bijective.

Example 5.35 *Following elements are minimal* $(0 \le l \le 7)$.

if $l \ge 0$ $\quad \phi,$

if $l \ge 1$ $\quad \Big\llbracket \llbracket 0_1 \rrbracket \Big\rrbracket,$

if $l \ge 2$ $\quad \Big\llbracket \llbracket 1 \rrbracket \llbracket \bar{1} \rrbracket \Big\rrbracket \quad \Big\llbracket \llbracket 3 \rrbracket \llbracket \bar{3} \rrbracket \Big\rrbracket,$

if $l \ge 3$ $\quad \Big\llbracket \llbracket 1 \rrbracket \llbracket 0_1 \rrbracket \llbracket \bar{1} \rrbracket \Big\rrbracket \quad \Big\llbracket \llbracket 2 \rrbracket \llbracket 0_2 \rrbracket \llbracket \bar{2} \rrbracket \Big\rrbracket,$

if $l \ge 4$ $\quad \Big\llbracket \llbracket 1 \rrbracket^2 \llbracket \bar{1} \rrbracket^2 \Big\rrbracket \quad \Big\llbracket \llbracket 1 \rrbracket \llbracket 3 \rrbracket \llbracket \bar{3} \rrbracket \llbracket \bar{1} \rrbracket \Big\rrbracket \quad \Big\llbracket \llbracket 2 \rrbracket \llbracket 4 \rrbracket \llbracket \bar{4} \rrbracket \llbracket \bar{2} \rrbracket \Big\rrbracket,$

if $l \ge 5$ $\quad \Big\llbracket \llbracket 1 \rrbracket^2 \llbracket 0_1 \rrbracket \llbracket \bar{1} \rrbracket^2 \Big\rrbracket \quad \Big\llbracket \llbracket 1 \rrbracket \llbracket 2 \rrbracket \llbracket 0_2 \rrbracket \llbracket \bar{2} \rrbracket \llbracket \bar{1} \rrbracket \Big\rrbracket \quad \Big\llbracket \llbracket 2 \rrbracket \llbracket 3 \rrbracket \llbracket 0_2 \rrbracket \llbracket \bar{3} \rrbracket \llbracket \bar{2} \rrbracket \Big\rrbracket,$

if $l \ge 6$ $\quad \Big\llbracket \llbracket 1 \rrbracket^3 \llbracket \bar{1} \rrbracket^3 \Big\rrbracket \quad \Big\llbracket \llbracket 1 \rrbracket^2 \llbracket 3 \rrbracket \llbracket \bar{3} \rrbracket \llbracket \bar{1} \rrbracket^2 \Big\rrbracket$

$\Big\llbracket \llbracket 1 \rrbracket \llbracket 2 \rrbracket \llbracket 4 \rrbracket \llbracket \bar{4} \rrbracket \llbracket \bar{2} \rrbracket \llbracket \bar{1} \rrbracket \Big\rrbracket \quad \Big\llbracket \llbracket 2 \rrbracket^2 \llbracket 6 \rrbracket \llbracket \bar{6} \rrbracket \llbracket \bar{2} \rrbracket^2 \Big\rrbracket,$

if $l \ge 7$ $\quad \Big\llbracket \llbracket 1 \rrbracket^3 \llbracket 0_1 \rrbracket \llbracket \bar{1} \rrbracket^3 \Big\rrbracket \quad \Big\llbracket \llbracket 1 \rrbracket^2 \llbracket 2 \rrbracket \llbracket 0_2 \rrbracket \llbracket \bar{2} \rrbracket \llbracket \bar{1} \rrbracket^2 \Big\rrbracket$

$\Big\llbracket \llbracket 1 \rrbracket \llbracket 2 \rrbracket \llbracket 3 \rrbracket \llbracket 0_2 \rrbracket \llbracket \bar{3} \rrbracket \llbracket \bar{2} \rrbracket \llbracket \bar{1} \rrbracket \Big\rrbracket \quad \Big\llbracket \llbracket 2 \rrbracket^2 \llbracket 4 \rrbracket \llbracket 0_2 \rrbracket \llbracket \bar{4} \rrbracket \llbracket \bar{2} \rrbracket^2 \Big\rrbracket.$

5.5 Connectedness of $B^l \otimes B^l$

We will show connectedness of $B^l \otimes B^l$, by showing that any element of $B^l \otimes B^l$ is connected with $\phi \otimes \phi$. We consider decomposition of tensor products $B^{G_2}(m_1\Lambda_1) \otimes B^{G_2}(m_2\Lambda_1)$. Each connected component has a lowest weight element. Then, we prove the lowest elements connected with $\phi \otimes \phi$.

For $b_1 \in B^{G_2}(m_1\Lambda_1)$, $b_2 \in B^{G_2}(m_2\Lambda_1)$, there exist a sequance $(i_1, \ldots, i_k)$ $(i_n \in \{1, 2\}, n = 1, \ldots, k)$ such that $\tilde{f}_{i_1} \cdots \tilde{f}_{i_2} b_1 \otimes b_2$ is a lowest weight element $b'_1 \otimes \Big\llbracket \llbracket \bar{1} \rrbracket^{m_2} \Big\rrbracket \in B^{G_2}(m_1\Lambda_1) \otimes B^{G_2}(m_2\Lambda_2)$. We can express

$$\tilde{f}_{i_1} \cdots \tilde{f}_{i_2} b_1 \otimes b_2 = b'_1 \otimes \Big\llbracket \llbracket \bar{1} \rrbracket^{m_2} \Big\rrbracket.$$

Then there exists $b''_1 \in B^{G_2}(m'_1\Lambda_1)$ such that

$$\tilde{f}_0^{(\varphi_0(b'_1) - m_2)_+ + m_2} b'_1 \otimes \Big\llbracket \llbracket \bar{1} \rrbracket^{m_2} \Big\rrbracket = b''_1 \otimes \phi.$$

There exist another sequence $(i'_1, \cdots, i'_{k'})$ $(i'_n \in \{1, 2\}, n = 1, \ldots, k')$ such that $\tilde{f}_0^{m'_1} \tilde{f}_{i'_1} \cdots \tilde{f}_{i'_{k'}} b''_1$ is lowest weight element $\Big\llbracket \llbracket \bar{1} \rrbracket^{m'_1} \Big\rrbracket$. Therefore we can express as

$$\tilde{f}_0^{m'_1} \tilde{f}_{i'_1} \cdots \tilde{f}_{i'_{k'}} b''_1 \otimes \phi = \tilde{f}_0^{m'_1} \Big\llbracket \llbracket \bar{1} \rrbracket^{m'_1} \Big\rrbracket \otimes \phi = \phi \otimes \phi.$$

Then we have

$$b_1 \otimes b_2 \text{ is connected with } \phi \otimes \phi.$$

By §5.4, §5.5 we have Theorem 2.2.

Perfect crystal of level 2


↙ $\tilde{f}_1$ ↘ $\tilde{f}_0$ other arrows: $E_{\mathcal{A}}$ $*$ is minimal.
$\mathcal{A}_0$
$\mathcal{A}_1$

Figure 9: Perfect crystal of level 2

References

[1] V.G.Drinfeld, Hopf algebra and the Yang-Baxter equation, *Soviet Math. Dokl.* **32** (1985), 254–258.

[2] M.Jimbo, A q-difference analogue of U($\mathfrak{g}$) and the Yang-Baxter equation, *Lett. Math. Phys.* **10** (1985),63–69.

[3] M.Kashiwara, Crystalizing the q-analogue of universal enveloping algebras, *Comm. Math. Phys.* **133** (1990), 249–260.

[4] M.Kashiwara, On crystal bases of the q-analogue of universal enveloping algebras, *Duke Math. J.* **63** (1991), 456–516.

[5] M.Kashiwara, Global crystal bases of quantum groups, *Duke Math. J.* **69** (1993), 455–485.

[6] M.Kashiwara and T.Nakashima, Crystal graphs for representations of q-analogue of classical Lie algebras, *J. Algebra* **165** (1994), 295–345.

[7] S-J.Kang and K.C.Misra, Crystal bases and tensor product decompositions of $U_q(G_2)$-module, *J. Algebra* **163** (1994), 675-691.

[8] S-J.Kang, M.Kashiwara, K.C.Misra, T.Miwa, T.Nakashima, and A.Nakayashiki, Perfect Crystal of Quantum Affine Lie Algebras, *Duke Math. J.* **68** (1992), 499–607.

[9] S-J.Kang, M.Kashiwara, K.C.Misra, T.Miwa, T.Nakashima, and A.Nakayashiki, Perfect Crystal and Vertex Models, *Internat. J. Modern Phys.* **A 7** (1992) 449–484.

[10] A.Kuniba, Quantum R Matrix for G_2 and a Solvable 175-vertex model, *J. Phys. A: Math. Gen.* **23** (1990) 1349–1362.

[11] P.Littelmann, A Littlewood-Richardson Rule for Symmetrizable Kac-Moody Algebra, *Invent. Math.* **116** (1994), 295–345.

[12] S-J.Kang, M.Kashiwara and K.C.Misra, Crystal bases of Verma Modules for Quantum Affine Lie Algebras, *Compositio Math* **92** (1994), 299-325.

[13] W.G.McKay, J.Patera and D.W.Rand TABLES OF REPRESENTATIONS OF SIMPLE LIE ALGEBLAS Volume I. Exceptional Simple Lie Algebras *Centre de Recherches Mathématiques Université de Montréal*